

1. Sugimoto K, Ikegami H, Takata Y, Katsuya T, Fukuda M, Akasaka H, Tabara Y, Osawa H, Hiromine Y, Rakugi H. Glycemic Control and Insulin Improve Muscle Mass and Gait Speed in Type 2 Diabetes: The MUSCLES-DM Study. *J Am Med Dir Assoc*. 2020;S1525-8610(20)30967-1.
2. Kario K, Hoshida S, Mizuno H, Kabutoya T, Nishizawa M, Yoshida T, Abe H, Katsuya T, Fujita Y, Okazaki O, Yano Y, Tomitani N, Kanegae H; JAMP Study Group. Nighttime Blood Pressure Phenotype and Cardiovascular Prognosis: Practitioner-Based Nationwide JAMP Study. 2020;142(19):1810-20.
3. Japan Geriatrics Society Subcommittee on End-of-Life Issues and New Coronavirus Countermeasure Team, Kuzuya M, Aita K, Katayama Y, Katsuya T, Nishikawa M, Hirahara S, Miura H, Yanagawa M, Arai H, Iijima K, Okochi J, Kozaki K, Yamaguchi Y, Rakugi H, Akishita M. The Japan Geriatrics Society consensus statement “recommendations for older persons to receive the best medical and long-term care during the COVID-19 outbreak-considering the timing of advance care planning implementation” *Gerontol. Int.* 2020 ; 20 (12) : 1112-19.
4. Japan Geriatrics Society Subcommittee on End-of-Life Issues, Kuzuya M, Aita K, Katayama Y, Katsuya T, Nishikawa M, Hirahara S, Miura H, Rakugi H, Akishita M. Japan Geriatrics Society “Recommendations for the Promotion of Advance Care Planning”: End-of-Life Issues Subcommittee consensus statement. *Gerontol. Int.* 2020 ; 20 (11) : 1024-28.
5. Spracklen CN, Horikoshi M, Kim YJ, Lin K, Bragg F, Moon S, Suzuki K, Tam CHT, Tabara Y, Kwak SH, Takeuchi F, Long J, Lim VJY, Chai JF, Chen CH, Nakatochi M, Yao J, Choi HS, Iyengar AK, Perrin HJ, Brotman SM, van de Bunt M, Gloyn AL, Below JE, Boehnke M, Bowden DW, Chambers JC, Mahajan A, McCarthy MI, Ng MCY, Petty LE, Zhang W, Morris AP, Adair LS, Akiyama M, Bian Z, Chan JCN, Chang LC, Chee ML, Chen YI, Chen YT, Chen Z, Chuang LM, Du S, Gordon-Larsen P, Gross M, Guo X, Guo Y, Han S, Howard AG, Huang W, Hung YJ, Hwang MY, Hwu CM, Ichihara S, Isono M, Jang HM, Jiang G, Jonas JB, Kamatani Y, Katsuya T, Kawaguchi T, Khor CC, Kohara K, Lee MS, Lee NR, Li L, Liu J, Luk AO, Lv J, Okada Y, Pereira MA, Sabanayagam C, Shi J, Shin DM, So WY, Takahashi A, Tomlinson B, Tsai FJ, van Dam RM, Xiang YB, Yamamoto K, Yamauchi T, Yoon K, Yu C, Yuan JM, Zhang L, Zheng W, Igase M, Cho YS, Rotter JI, Wang YX, Sheu WHH, Yokota M, Wu JY, Cheng CY, Wong TY, Shu XO, Kato N, Park KS, Tai ES, Matsuda F, Koh WP, Ma RCW, Maeda S, Millwood IY, Lee J, Kadowaki T, Walters RG, Kim BJ, Mohlke KL, Sim X. Identification of type 2 diabetes loci in 433,540 East Asian individuals. *Nature*. 2020;582(7811):240-245.
6. Kasahara M, Kuwabara Y, Moriyama T, Tanabe K, Satoh-Asahara N, Katsuya T, Hiramitsu S, Shimada H, Sato T, Saito Y, Nakagawa T. Intensive uric acid-lowering therapy in CKD patients: the protocol for a randomized controlled trial. *Clin Exp Nephrol*. 2020;24:235-241.
7. Asayama K, Ohkubo T, Rakugi H, Miyakawa M, Mori H, Katsuya T, Ikehara Y, Ueda S, Ohya Y, Tsuchihashi T, Kario K, Miura K, Hasebe N, Ito S, Umemura S; Japanese Society of Hypertension

Working Group on the COmparison of Self-measured home, Automated unattended office and Conventional attended office blood pressure (COSAC) study. Comparison of blood pressure values-self-measured at home, measured at an unattended office, and measured at a conventional attended office. *Hypertens Res.* 2019;42(11):1726-37.

8. Sung YJ, de Las Fuentes L, Winkler TW, Chasman DI, Bentley AR, Kraja AT, Ntalla I, Warren HR, Guo X, Schwander K, Manning AK, Brown MR, Aschard H, Feitosa MF, Franceschini N, Lu Y, Cheng CY, Sim X, Vojinovic D, Marten J, Musani SK, Kilpeläinen TO, Richard MA, Aslibekyan S, Bartz TM, Dorajoo R, Li C, Liu Y, Rankinen T, Smith AV, Tajuddin SM, Tayo BO, Zhao W, Zhou Y, Matoba N, Sofer T, Alver M, Amini M, Boissel M, Chai JF, Chen X, Divers J, Gandin I, Gao C, Giulianini F, Goel A, Harris SE, Hartwig FP, He M, Horimoto ARVR, Hsu FC, Jackson AU, Kammerer CM, Kasturiratne A, Komulainen P, Kühnel B, Leander K, Lee WJ, Lin KH, Luan J, Lyytikäinen LP, McKenzie CA, Nelson CP, Noordam R, Scott RA, Sheu WHH, Stančáková A, Takeuchi F, van der Most PJ, Varga TV, Waken RJ, Wang H, Wang Y, Ware EB, Weiss S, Wen W, Yanek LR, Zhang W, Zhao JH, Afaq S, Alfred T, Amin N, Arking DE, Aung T, Barr RG, Bielak LF, Boerwinkle E, Bottinger EP, Braund PS, Brody JA, Broeckel U, Cade B, Campbell A, Canouil M, Chakravarti A, Cocca M, Collins FS, Connell JM, de Mutsert R, de Silva HJ, Dörr M, Duan Q, Eaton CB, Ehret G, Evangelou E, Faul JD, Forouhi NG, Franco OH, Friedlander Y, Gao H, Gigante B, Gu CC, Gupta P, Hagenaars SP, Harris TB, He J, Heikkinen S, Heng CK, Hofman A, Howard BV, Hunt SC, Irvin MR, Jia Y, Katsuya T, Kaufman J, Kerrison ND, Khor CC, Koh WP, Koistinen HA, Kooperberg CB, Krieger JE, Kubo M, Kutalik Z, Kuusisto J, Lakka TA, Langefeld CD, Langenberg C, Launer LJ, Lee JH, Lehne B, Levy D, Lewis CE, Li Y; Lifelines Cohort Study, Lim SH, Liu CT, Liu J, Liu J, Liu Y, Loh M, Lohman KK, Louie T, Mägi R, Matsuda K, Meitinger T, Metspalu A, Milani L, Momozawa Y, Mosley TH Jr, Nalls MA, Nasri U, O'Connell JR, Ogunniyi A, Palmas WR, Palmer ND, Pankow JS, Pedersen NL, Peters A, Peyser PA, Polasek O, Porteous D, Raitakari OT, Renström F, Rice TK, Ridker PM, Robino A, Robinson JG, Rose LM, Rudan I, Sabanayagam C, Salako BL, Sandow K, Schmidt CO, Schreiner PJ, Scott WR, Sever P, Sims M, Sitlani CM, Smith BH, Smith JA, Snieder H, Starr JM, Strauch K, Tang H, Taylor KD, Teo YY, Tham YC, Uitterlinden AG, Waldenberger M, Wang L, Wang YX, Wei WB, Wilson G, Wojczynski MK, Xiang YB, Yao J, Yuan JM, Zonderman AB, Becker DM, Boehnke M, Bowden DW, Chambers JC, Chen YI, Weir DR, de Faire U, Deary IJ, Esko T, Farrall M, Forrester T, Freedman BI, Froguel P, Gasparini P, Gieger C, Horta BL, Hung YJ, Jonas JB, Kato N, Kooner JS, Laakso M, Lehtimäki T, Liang KW, Magnusson PKE, Oldehinkel AJ, Pereira AC, Perls T, Rauramaa R, Redline S, Rettig R, Samani NJ, Scott J, Shu XO, van der Harst P, Wagenknecht LE, Wareham NJ, Watkins H, Wickremasinghe AR, Wu T, Kamatani Y, Laurie CC, Bouchard C, Cooper RS, Evans MK, Gudnason V, Hixson J, Kardia SLR, Kritchevsky SB, Psaty BM, van Dam RM, Arnett DK, Mook-Kanamori DO, Fornage M, Fox ER, Hayward C, van Duijn CM, Tai ES, Wong TY, Loos RJJ, Reiner AP, Rotimi CN, Bierut LJ, Zhu X, Cupples LA, Province MA, Rotter JI, Franks PW, Rice K, Elliott P, Caulfield MJ, Gauderman WJ, Munroe PB, Rao DC, Morrison

AC. A multi-ancestry genome-wide study incorporating gene-smoking interactions identifies multiple new loci for pulse pressure and mean arterial pressure. *Hum Mol Genet.* 2019;28(15):2615-33.

9. Sugimoto K, Tabara Y, Ikegami H, Takata Y, Kamide K, Ikezoe T, Kiyoshige E, Makutani Y, Onuma H, Gondo Y, Ikebe K, Ichihashi N, Tsuboyama T, Matsuda F, Kohara K, Kabayama M, Fukuda M, Katsuya T, Osawa H, Hiromine Y, Rakugi H. Hyperglycemia in non-obese patients with type 2 diabetes is associated with low muscle mass: The MUSCLES-DM study. *J Diabetes Investig.* 2019;10(6):1471-9.
10. Bentley AR, Sung YJ, Brown MR, Winkler TW, Kraja AT, Ntalla I, Schwander K, Chasman DI, Lim E, Deng X, Guo X, Liu J, Lu Y, Cheng CY, Sim X, Vojinovic D, Huffman JE, Musani SK, Li C, Feitosa MF, Richard MA, Noordam R, Baker J, Chen G, Aschard H, Bartz TM, Ding J, Dorajoo R, Manning AK, Rankinen T, Smith AV, Tajuddin SM, Zhao W, Graff M, Alver M, Boissel M, Chai JF, Chen X, Divers J, Evangelou E, Gao C, Goel A, Hagemeijer Y, Harris SE, Hartwig FP, He M, Horimoto ARVR, Hsu FC, Hung YJ, Jackson AU, Kasturiratne A, Komulainen P, Kühnel B, Leander K, Lin KH, Luan J, Lyytikäinen LP, Matoba N, Nolte IM, Pietzner M, Prins B, Riaz M, Robino A, Said MA, Schupf N, Scott RA, Sofer T, Stancáková A, Takeuchi F, Tayo BO, van der Most PJ, Varga TV, Wang TD, Wang Y, Ware EB, Wen W, Xiang YB, Yanek LR, Zhang W, Zhao JH, Adeyemo A, Afaq S, Amin N, Amini M, Arking DE, Arzumanyan Z, Aung T, Ballantyne C, Barr RG, Bielak LF, Boerwinkle E, Bottinger EP, Broeckel U, Brown M, Cade BE, Campbell A, Canouil M, Charumathi S, Chen YI, Christensen K; COGENT-Kidney Consortium, Concas MP, Connell JM, de Las Fuentes L, de Silva HJ, de Vries PS, Doumatey A, Duan Q, Eaton CB, Eppinga RN, Faul JD, Floyd JS, Forouhi NG, Forrester T, Friedlander Y, Gandin I, Gao H, Ghanbari M, Gharib SA, Gigante B, Giulianini F, Grabe HJ, Gu CC, Harris TB, Heikkinen S, Heng CK, Hirata M, Hixson JE, Ikram MA; EPIC-InterAct Consortium, Jia Y, Joeannes R, Johnson C, Jonas JB, Justice AE, Katsuya T, Khor CC, Kilpeläinen TO, Koh WP, Kolcic I, Kooperberg C, Krieger JE, Kritchevsky SB, Kubo M, Kuusisto J, Lakka TA, Langefeld CD, Langenberg C, Launer LJ, Lehne B, Lewis CE, Li Y, Liang J, Lin S, Liu CT, Liu J, Liu K, Loh M, Lohman KK, Louie T, Luzzi A, Mägi R, Mahajan A, Manichaikul AW, McKenzie CA, Meitinger T, Metspalu A, Milaneschi Y, Milani L, Mohlke KL, Momozawa Y, Morris AP, Murray AD, Nalls MA, Nauck M, Nelson CP, North KE, O'Connell JR, Palmer ND, Papanicolaou GJ, Pedersen NL, Peters A, Peyser PA, Polasek O, Poulter N, Raitakari OT, Reiner AP, Renström F, Rice TK, Rich SS, Robinson JG, Rose LM, Rosendaal FR, Rudan I, Schmidt CO, Schreiner PJ, Scott WR, Sever P, Shi Y, Sidney S, Sims M, Smith JA, Snieder H, Starr JM, Strauch K, Stringham HM, Tan NYQ, Tang H, Taylor KD, Teo YY, Tham YC, Tiemeier H, Turner ST, Uitterlinden AG; Understanding Society Scientific Group, van Heemst D, Waldenberger M, Wang H, Wang L, Wang L, Wei WB, Williams CA, Wilson G Sr, Wojczynski MK, Yao J, Young K, Yu C, Yuan JM, Zhou J, Zonderman AB, Becker DM, Boehnke M, Bowden DW, Chambers JC, Cooper RS, de Faire U, Deary IJ, Elliott P, Esko T, Farrall M, Franks PW, Freedman BI, Froguel P, Gasparini P, Gieger C, Horta BL, Juang JJ, Kamatani Y, Kammerer CM, Kato N, Kooner JS, Laakso M, Laurie CC, Lee IT, Lehtimäki T;

Lifelines Cohort, Magnusson PKE, Oldehinkel AJ, Penninx BWJH, Pereira AC, Rauramaa R, Redline S, Samani NJ, Scott J, Shu XO, van der Harst P, Wagenknecht LE, Wang JS, Wang YX, Wareham NJ, Watkins H, Weir DR, Wickremasinghe AR, Wu T, Zeggini E, Zheng W, Bouchard C, Evans MK, Gudnason V, Kardia SLR, Liu Y, Psaty BM, Ridker PM, van Dam RM, Mook-Kanamori DO, Fornage M, Province MA, Kelly TN, Fox ER, Hayward C, van Duijn CM, Tai ES, Wong TY, Loos RJF, Franceschini N, Rotter JI, Zhu X, Bierut LJ, Gauderman WJ, Rice K, Munroe PB, Morrison AC, Rao DC, Rotimi CN, Cupples LA. Multi-ancestry genome-wide gene-smoking interaction study of 387,272 individuals identifies new loci associated with serum lipids. *Nat Genet.* 2019;51(4):636-648.

11. de Vries PS, Brown MR, Bentley AR, Sung YJ, Winkler TW, Ntalla I, Schwander K, Kraja AT, Guo X, Franceschini N, Cheng CY, Sim X, Vojinovic D, Huffman JE, Musani SK, Li C, Feitosa MF, Richard MA, Noordam R, Aschard H, Bartz TM, Bielak LF, Deng X, Dorajoo R, Lohman KK, Manning AK, Rankinen T, Smith AV, Tajuddin SM, Evangelou E, Graff M, Alver M, Boissel M, Chai JF, Chen X, Divers J, Gandin I, Gao C, Goel A, Hagemeijer Y, Harris SE, Hartwig FP, He M, Horimoto ARVR, Hsu FC, Jackson AU, Kasturiratne A, Komulainen P, Kühnel B, Laguzzi F, Lee JH, Luan J, Lyytikäinen LP, Matoba N, Nolte IM, Pietzner M, Riaz M, Said MA, Scott RA, Sofer T, Stancáková A, Takeuchi F, Tayo BO, van der Most PJ, Varga TV, Wang Y, Ware EB, Wen W, Yanek LR, Zhang W, Zhao JH, Afaq S, Amin N, Amini M, Arking DE, Aung T, Ballantyne C, Boerwinkle E, Broeckel U, Campbell A, Canouil M, Charumathi S, Chen YI, Connell JM, de Faire U, de Las Fuentes L, de Mutsert R, de Silva HJ, Ding J, Dominiczak AF, Duan Q, Eaton CB, Eppinga RN, Faul JD, Fisher V, Forrester T, Franco OH, Friedlander Y, Ghanbari M, Giulianini F, Grabe HJ, Grove ML, Gu CC, Harris TB, Heikkinen S, Heng CK, Hirata M, Hixson JE, Howard BV, Ikram MA; InterAct Consortium, Jacobs DR Jr, Johnson C, Jonas JB, Kammerer CM, Katsuya T, Khor CC, Kilpeläinen TO, Koh WP, Koistinen HA, Kolcic I, Kooperberg C, Krieger JE, Kritchevsky SB, Kubo M, Kuusisto J, Lakka TA, Langeveld CD, Langenberg C, Launer LJ, Lehne B, Lemaitre RN, Li Y, Liang J, Liu J, Liu K, Loh M, Louie T, Mägi R, Manichaikul AW, McKenzie CA, Meitinger T, Metspalu A, Milaneschi Y, Milani L, Mohlke KL, Mosley TH Jr, Mukamal KJ, Nalls MA, Nauck M, Nelson CP, Sotoodehnia N, O'Connell JR, Palmer ND, Pazoki R, Pedersen NL, Peters A, Peyser PA, Polasek O, Poulter N, Raffel LJ, Raitakari OT, Reiner AP, Rice TK, Rich SS, Robino A, Robinson JG, Rose LM, Rudan I, Schmidt CO, Schreiner PJ, Scott WR, Sever P, Shi Y, Sidney S, Sims M, Smith BH, Smith JA, Snieder H, Starr JM, Strauch K, Tan N, Taylor KD, Teo YY, Tham YC, Uitterlinden AG, van Heemst D, Vuckovic D, Waldenberger M, Wang L, Wang Y, Wang Z, Wei WB, Williams C, Wilson G Sr, Wojczynski MK, Yao J, Yu B, Yu C, Yuan JM, Zhao W, Zonderman AB, Becker DM, Boehnke M, Bowden DW, Chambers JC, Deary IJ, Esko T, Farrall M, Franks PW, Freedman BI, Froguel P, Gasparini P, Gieger C, Horta BL, Kamatani Y, Kato N, Kooner JS, Laakso M, Leander K, Lehtimäki T; Lifelines Cohort, Groningen, The Netherlands (Lifelines Cohort Study), Magnusson PKE, Penninx B, Pereira AC, Rauramaa R, Samani NJ, Scott J, Shu XO, van der Harst P, Wagenknecht LE, Wang YX, Wareham NJ, Watkins H, Weir DR, Wickremasinghe AR, Zheng W, Elliott P, North KE, Bouchard C,

Evans MK, Gudnason V, Liu CT, Liu Y, Psaty BM, Ridker PM, van Dam RM, Kardia SLR, Zhu X, Rotimi CN, Mook-Kanamori DO, Fornage M, Kelly TN, Fox ER, Hayward C, van Duijn CM, Tai ES, Wong TY, Liu J, Rotter JI, Gauderman WJ, Province MA, Munroe PB, Rice K, Chasman DI, Cupples LA, Rao DC, Morrison AC. Multi-Ancestry Genome-Wide Association Study of Lipid Levels Incorporating Gene-Alcohol Interactions. *Am J Epidemiol*. 2019;188(6):1033-54.

12. Kario K, Okada K, Kato M, Nishizawa M, Yoshida T, Asano T, Uchiyama K, Niiijima Y, Katsuya T, Urata H, Osuga JI, Fujiwara T, Yamazaki S, Tomitani N, Kanegae H. 24-Hour Blood Pressure-Lowering Effect of an SGLT-2 Inhibitor in Patients with Diabetes and Uncontrolled Nocturnal Hypertension: Results from the Randomized, Placebo-Controlled SACRA Study. *Circulation*. 2019;139:2089-97.

13. Takeuchi F, Akiyama M, Matoba N, Katsuya T, Nakatochi M, Tabara Y, Narita A, Saw WY, Moon S, Spracklen CN, Chai JF, Kim YJ, Zhang L, Wang C, Li H, Li H, Wu JY, Dorajoo R, Nierenberg JL, Wang YX, He J, Bennett DA, Takahashi A, Momozawa Y, Hirata M, Matsuda K, Rakugi H, Nakashima E, Isono M, Shirota M, Hozawa A, Ichihara S, Matsubara T, Yamamoto K, Kohara K, Igase M, Han S, Gordon-Larsen P, Huang W, Lee NR, Adair LS, Hwang MY, Lee J, Chee ML, Sabanayagam C, Zhao W, Liu J, Reilly DF, Sun L, Huo S, Edwards TL, Long J, Chang LC, Chen CH, Yuan JM, Koh WP, Friedlander Y, Kelly TN, Bin Wei W, Xu L, Cai H, Xiang YB, Lin K, Clarke R, Walters RG, Millwood IY, Li L, Chambers JC, Kooner JS, Elliott P, van der Harst P; International Genomics of Blood Pressure (iGEN-BP) Consortium, Chen Z, Sasaki M, Shu XO, Jonas JB, He J, Heng CK, Chen YT, Zheng W, Lin X, Teo YY, Tai ES, Cheng CY, Wong TY, Sim X, Mohlke KL, Yamamoto M, Kim BJ, Miki T, Nabika T, Yokota M, Kamatani Y, Kubo M, Kato N. Interethnic analyses of blood pressure loci in populations of East Asian and European descent. *Nat Commun*. 2018;9(1):5052.

14. Feitosa MF, Kraja AT, Chasman DI, Sung YJ, Winkler TW, Ntalla I, Guo X, Franceschini N, Cheng CY, Sim X, Vojinovic D, Marten J, Musani SK, Li C, Bentley AR, Brown MR, Schwander K, Richard MA, Noordam R, Aschard H, Bartz TM, Bielak LF, Dorajoo R, Fisher V, Hartwig FP, Horimoto ARVR, Lohman KK, Manning AK, Rankinen T, Smith AV, Tajuddin SM, Wojczynski MK, Alver M, Boissel M, Cai Q, Campbell A, Chai JF, Chen X, Divers J, Gao C, Goel A, Hagemeijer Y, Harris SE, He M, Hsu FC, Jackson AU, Kähönen M, Kasturiratne A, Komulainen P, Kühnel B, Laguzzi F, Luan J, Matoba N, Nolte IM, Padmanabhan S, Riaz M, Rueedi R, Robino A, Said MA, Scott RA, Sofer T, Stančáková A, Takeuchi F, Tayo BO, van der Most PJ, Varga TV, Vitart V, Wang Y, Ware EB, Warren HR, Weiss S, Wen W, Yanek LR, Zhang W, Zhao JH, Afaq S, Amin N, Amini M, Arking DE, Aung T, Boerwinkle E, Borecki I, Broeckel U, Brown M, Brumat M, Burke GL, Canouil M, Chakravarti A, Charumathi S, Ida Chen YD, Connell JM, Correa A, de Las Fuentes L, de Mutsert R, de Silva HJ, Deng X, Ding J, Duan Q, Eaton CB, Ehret G, Eppinga RN, Evangelou E, Faul JD, Felix SB, Forouhi NG, Forrester T, Franco OH, Friedlander Y, Gandin I, Gao H, Ghanbari M, Gigante B, Gu CC, Gu D, Hagenaars SP, Hallmans G, Harris TB, He J, Heikkinen S, Heng CK, Hirata M, Howard BV, Ikram MA; InterAct Consortium, John U, Katsuya T,

Khor CC, Kilpeläinen TO, Koh WP, Krieger JE, Kritchevsky SB, Kubo M, Kuusisto J, Lakka TA, Langefeld CD, Langenberg C, Launer LJ, Lehne B, Lewis CE, Li Y, Lin S, Liu J, Liu J, Loh M, Louie T, Mägi R, McKenzie CA, Meitinger T, Metspalu A, Milaneschi Y, Milani L, Mohlke KL, Momozawa Y, Nalls MA, Nelson CP, Sotoodehnia N, Norris JM, O'Connell JR, Palmer ND, Perls T, Pedersen NL, Peters A, Peyser PA, Poulter N, Raffel LJ, Raitakari OT, Roll K, Rose LM, Rosendaal FR, Rotter JI, Schmidt CO, Schreiner PJ, Schupf N, Scott WR, Sever PS, Shi Y, Sidney S, Sims M, Sitlani CM, Smith JA, Snieder H, Starr JM, Strauch K, Stringham HM, Tan NYQ, Tang H, Taylor KD, Teo YY, Tham YC, Turner ST, Uitterlinden AG, Vollenweider P, Waldenberger M, Wang L, Wang YX, Wei WB, Williams C, Yao J, Yu C, Yuan JM, Zhao W, Zonderman AB, Becker DM, Boehnke M, Bowden DW, Chambers JC, Deary IJ, Esko T, Farrall M, Franks PW, Freedman BI, Froguel P, Gasparini P, Gieger C, Jonas JB, Kamatani Y, Kato N, Kooner JS, Kutalik Z, Laakso M, Laurie CC, Leander K, Lehtimäki T, Study LC, Magnusson PKE, Oldehinkel AJ, Penninx BWJH, Polasek O, Porteous DJ, Rauramaa R, Samani NJ, Scott J, Shu XO, van der Harst P, Wagenknecht LE, Wareham NJ, Watkins H, Weir DR, Wickremasinghe AR, Wu T, Zheng W, Bouchard C, Christensen K, Evans MK, Gudnason V, Horta BL, Kardia SLR, Liu Y, Pereira AC, Psaty BM, Ridker PM, van Dam RM, Gauderman WJ, Zhu X, Mook-Kanamori DO, Fornage M, Rotimi CN, Cupples LA, Kelly TN, Fox ER, Hayward C, van Duijn CM, Tai ES, Wong TY, Kooperberg C, Palmas W, Rice K, Morrison AC, Elliott P, Caulfield MJ, Munroe PB, Rao DC, Province MA, Levy D. Novel genetic associations for blood pressure identified via gene-alcohol interaction in up to 570K individuals across multiple ancestries. *PLoS One*. 2018;13(6):e0198166.

15. Sung YJ, Winkler TW, de Las Fuentes L, Bentley AR, Brown MR, Kraja AT, Schwander K, Ntalla I, Guo X, Franceschini N, Lu Y, Cheng CY, Sim X, Vojinovic D, Marten J, Musani SK, Li C, Feitosa MF, Kilpeläinen TO, Richard MA, Noordam R, Aslibekyan S, Aschard H, Bartz TM, Dorajoo R, Liu Y, Manning AK, Rankinen T, Smith AV, Tajuddin SM, Tayo BO, Warren HR, Zhao W, Zhou Y, Matoba N, Sofer T, Alver M, Amini M, Boissel M, Chai JF, Chen X, Divers J, Gandin I, Gao C, Giulianini F, Goel A, Harris SE, Hartwig FP, Horimoto ARVR, Hsu FC, Jackson AU, Kähönen M, Kasturiratne A, Kühnel B, Leander K, Lee WJ, Lin KH, 'an Luan J, McKenzie CA, Meian H, Nelson CP, Rauramaa R, Schupf N, Scott RA, Sheu WHH, Stančáková A, Takeuchi F, van der Most PJ, Varga TV, Wang H, Wang Y, Ware EB, Weiss S, Wen W, Yanek LR, Zhang W, Zhao JH, Afaq S, Alfred T, Amin N, Arking D, Aung T, Barr RG, Bielak LF, Boerwinkle E, Bottinger EP, Braund PS, Brody JA, Broeckel U, Cabrera CP, Cade B, Caizheng Y, Campbell A, Canouil M, Chakravarti A; CHARGE Neurology Working Group, Chauhan G, Christensen K, Cocca M; COGENT-Kidney Consortium, Collins FS, Connell JM, de Mutsert R, de Silva HJ, Debette S, Dörr M, Duan Q, Eaton CB, Ehret G, Evangelou E, Faul JD, Fisher VA, Forouhi NG, Franco OH, Friedlander Y, Gao H; GIANT Consortium, Gigante B, Graff M, Gu CC, Gu D, Gupta P, Hagenaars SP, Harris TB, He J, Heikkinen S, Heng CK, Hirata M, Hofman A, Howard BV, Hunt S, Irvin MR, Jia Y, Joehanes R, Justice AE, Katsuya T, Kaufman J, Kerrison ND, Khor CC, Koh WP, Koistinen HA, Komulainen P, Kooperberg C, Krieger JE, Kubo M, Kuusisto J, Langefeld CD, Langenberg C,

- Launer LJ, Lehne B, Lewis CE, Li Y; Lifelines Cohort Study, Lim SH, Lin S, Liu CT, Liu J, Liu J, Liu K, Liu Y, Loh M, Lohman KK, Long J, Louie T, Mägi R, Mahajan A, Meitinger T, Metspalu A, Milani L, Momozawa Y, Morris AP, Mosley TH Jr., Munson P, Murray AD, Nalls MA, Nasri U, Norris JM, North K, Ogunniyi A, Padmanabhan S, Palmas WR, Palmer ND, Pankow JS, Pedersen NL, Peters A, Peyser PA, Polasek O, Raitakari OT, Renström F, Rice TK, Ridker PM, Robino A, Robinson JG, Rose LM, Rudan I, Sabanayagam C, Salako BL, Sandow K, Schmidt CO, Schreiner PJ, Scott WR, Seshadri S, Sever P, Sitlani CM, Smith JA, Snieder H, Starr JM, Strauch K, Tang H, Taylor KD, Teo YY, Tham YC, Uitterlinden AG, Waldenberger M, Wang L, Wang YX, Wei WB, Williams C, Wilson G, Wojczynski MK, Yao J, Yuan JM, Zonderman AB, Becker DM, Boehnke M, Bowden DW, Chambers JC, Chen YI, de Faire U, Deary IJ, Esko T, Farrall M, Forrester T, Franks PW, Freedman BI, Froguel P, Gasparini P, Gieger C, Horta BL, Hung YJ, Jonas JB, Kato N, Kooner JS, Laakso M, Lehtimäki T, Liang KW, Magnusson PKE, Newman AB, Oldehinkel AJ, Pereira AC, Redline S, Rettig R, Samani NJ, Scott J, Shu XO, van der Harst P, Wagenknecht LE, Wareham NJ, Watkins H, Weir DR, Wickremasinghe AR, Wu T, Zheng W, Kamatani Y, Laurie CC, Bouchard C, Cooper RS, Evans MK, Gudnason V, Kardia SLR, Kritchevsky SB, Levy D, O'Connell JR, Psaty BM, van Dam RM, Sims M, Arnett DK, Mook-Kanamori DO, Kelly TN, Fox ER, Hayward C, Fornage M, Rotimi CN, Province MA, van Duijn CM, Tai ES, Wong TY, Loos RJF, Reiner AP, Rotter JI, Zhu X, Bierut LJ, Gauderman WJ, Caulfield MJ, Elliott P, Rice K, Munroe PB, Morrison AC, Cupples LA, Rao DC, Chasman DI. A large-scale multi-ancestry genome-wide study accounting for smoking behavior identifies multiple significant loci for blood pressure. *Am J Hum Genet* 2018;102(3):375-400.
- 16.Ogata S, Kamide K, Asayama K, Tabara Y, Kawaguchi T, Satoh M, Katsuya T, Sugimoto K, Hirose T, Inoue R, Hara A, Obara T, Kikuya M, Metoki H, Matsuda F, Staessen JA, Ohkubo T, Rakugi H, Imai Y. Genome-wide association study for white coat effect in Japanese middle-aged to elderly people: The HOMED-BP study. *Clin Exp Hypertens*. 2018;40(4):363-9.
- 17.Takeuchi F, Katsuya T, Kimura R, Nabika T, Isomura M, Ohkubo T, Tabara Y, Yamamoto K, Yokota M, Liu X, Saw WY, Mamatyusupu D, Yang W, Xu S; Japanese Genome Variation Consortium, Teo YY, Kato N. The fine-scale genetic structure and evolution of the Japanese population. *PLoS One*. 2017;12(11):e0185487.
- 18.Wheeler E, Leong A, Liu CT, Hivert MF, Strawbridge RJ, Podmore C, Li M, Yao J, Sim X, Hong J, Chu AY, Zhang W, Wang X, Chen P, Maruthur NM, Porneala BC, Sharp SJ, Jia Y, Kabagambe EK, Chang LC, Chen WM, Elks CE, Evans DS, Fan Q, Giulianini F, Go MJ, Hottenga JJ, Hu Y, Jackson AU, Kanoni S, Kim YJ, Kleber ME, Ladenvall C, Lecoeur C, Lim SH, Lu Y, Mahajan A, Marzi C, Nalls MA, Navarro P, Nolte IM, Rose LM, Rybin DV, Sanna S, Shi Y, Stram DO, Takeuchi F, Tan SP, van der Most PJ, Van Vliet-Ostaptchouk JV, Wong A, Yengo L, Zhao W, Goel A, Martinez Larrad MT, Radke D, Salo P, Tanaka T, van Iperen EPA, Abecasis G, Afaq S, Alizadeh BZ, Bertoni AG, Bonnefond A, Böttcher Y,

Bottinger EP, Campbell H, Carlson OD, Chen CH, Cho YS, Garvey WT, Gieger C, Goodarzi MO, Grallert H, Hamsten A, Hartman CA, Herder C, Hsiung CA, Huang J, Igase M, Isono M, Katsuya T, Khor CC, Kiess W, Kohara K, Kovacs P, Lee J, Lee WJ, Lehne B, Li H, Liu J, Lobbens S, Luan J, Lyssenko V, Meitinger T, Miki T, Miljkovic I, Moon S, Mulas A, Müller G, Müller-Nurasyid M, Nagaraja R, Nauck M, Pankow JS, Polasek O, Prokopenko I, Ramos PS, Rasmussen-Torvik L, Rathmann W, Rich SS, Robertson NR, Roden M, Roussel R, Rudan I, Scott RA, Scott WR, Sennblad B, Siscovick DS, Strauch K, Sun L, Swertz M, Tajuddin SM, Taylor KD, Teo YY, Tham YC, Tönjes A, Wareham NJ, Willemssen G, Wilsgaard T, Hingorani AD; EPIC-CVD Consortium; EPIC-InterAct Consortium; Lifelines Cohort Study, Egan J, Ferrucci L, Hovingh GK, Jula A, Kivimaki M, Kumari M, Njølstad I, Palmer CNA, Serrano Ríos M, Stumvoll M, Watkins H, Aung T, Blüher M, Boehnke M, Boomsma DI, Bornstein SR, Chambers JC, Chasman DI, Chen YI, Chen YT, Cheng CY, Cucca F, de Geus EJC, Deloukas P, Evans MK, Fornage M, Friedlander Y, Froguel P, Groop L, Gross MD, Harris TB, Hayward C, Heng CK, Ingelsson E, Kato N, Kim BJ, Koh WP, Kooner JS, Körner A, Kuh D, Kuusisto J, Laakso M, Lin X, Liu Y, Loos RJF, Magnusson PKE, März W, McCarthy MI, Oldehinkel AJ, Ong KK, Pedersen NL, Pereira MA, Peters A, Ridker PM, Sabanayagam C, Sale M, Saleheen D, Saltevo J, Schwarz PE, Sheu WHH, Snieder H, Spector TD, Tabara Y, Tuomilehto J, van Dam RM, Wilson JG, Wilson JF, Wolffenbuttel BHR, Wong TY, Wu JY, Yuan JM, Zonderman AB, Soranzo N, Guo X, Roberts DJ, Florez JC, Sladek R, Dupuis J, Morris AP, Tai ES, Selvin E, Rotter JI, Langenberg C, Barroso I, Meigs JB. Impact of common genetic determinants of Hemoglobin A1c on type 2 diabetes risk and diagnosis in ancestrally diverse populations: A transethnic genome-wide meta-analysis. *PLoS Med.* 2017 Sep 12;14(9):e1002383.

19. Nomura A, Won HH, Khera AV, Takeuchi F, Ito K, McCarthy S, Emdin CA, Klarin D, Natarajan P, Zekavat SM, Gupta N, Peloso GM, Borecki IB, Teslovich TM, Asselta R, Duga S, Merlini PA, Correa A, Kessler T, Wilson JG, Bown MJ, Hall AS, Braund PS, Carey DJ, Murray MF, Kirchner HL, Leader JB, Lavage DR, Manus JN, Hartzel DN, Samani NJ, Schunkert H, Marrugat J, Elosua R, McPherson R, Farrall M, Watkins H, Juang JJ, Hsiung CA, Lin SY, Wang JS, Tada H, Kawashiri MA, Inazu A, Yamagishi M, Katsuya T, Nakashima E, Nakatochi M, Yamamoto K, Yokota M, Momozawa Y, Rotter JI, Lander ES, Rader DJ, Danesh JN, Ardisson D, Gabriel SB, Willer CJ, Abecasis G, Saleheen D, Kubo M, Kato N, Chen YI, Dewey F, Kathiresan S. Protein truncating variants at the cholesteryl ester transfer protein gene and risk for coronary heart disease. *Circ Res.* 2017;121:81-8.

20. Kario K, Tomitani N, Kanegae H, Ishii H, Uchiyama K, Yamagiwa K, Shiraiwa T, Katsuya T, Yoshida T, Kanda K, Hasegawa S, Hoshida S. Comparative Effects of an Angiotensin II Receptor Blocker (ARB)/Diuretic vs. ARB/Calcium-Channel Blocker Combination on Uncontrolled Nocturnal Hypertension Evaluated by Information and Communication Technology-Based Nocturnal Home Blood Pressure Monitoring - The NOCTURNE Study. *Circ J.* 2017;81:948-57.

21. Spracklen CN, Chen P, Kim YJ, Wang X, Cai H, Li S, Long J, Wu Y, Wang YX, Takeuchi F, Wu JY, Jung KJ, Hu C, Akiyama K, Zhang Y, Moon S, Johnson TA, Li H, Dorajoo R, He M, Cannon ME, Roman TS, Salfati E, Lin KH, Guo X, Sheu WH, Absher D, Adair LS, Assimes TL, Aung T, Cai Q, Chang LC, Chen CH, Chien LH, Chuang LM, Chuang SC, Du S, Fan Q, Fann CS, Feranil AB, Friedlander Y, Gordon-Larsen P, Gu D, Gui L, Guo Z, Heng CK, Hixson J, Hou X, Hsiung CA, Hu Y, Hwang MY, Hwu CM, Isono M, Juang JJ, Khor CC, Kim YK, Koh WP, Kubo M, Lee IT, Lee SJ, Lee WJ, Liang KW, Lim B, Lim SH, Liu J, Nabika T, Pan WH, Peng H, Quertermous T, Sabanayagam C, Sandow K, Shi J, Sun L, Tan PC, Tan SP, Taylor KD, Teo YY, Toh SA, Tsunoda T, van Dam RM, Wang A, Wang F, Wang J, Wei WB, Xiang YB, Yao J, Yuan JM, Zhang R, Zhao W, Chen YI, Rich SS, Rotter JI, Wang TD, Wu T, Lin X, Han BG, Tanaka T, Cho YS, Katsuya T, Jia W, Jee SH, Chen YT, Kato N, Jonas JB, Cheng CY, Shu XO, He J, Zheng W, Wong TY, Huang W, Kim BJ, Tai ES, Mohlke KL, Sim X. Association analyses of East Asian individuals and trans-ancestry analyses with European individuals reveal new loci associated with cholesterol and triglyceride levels. *Hum Mol Genet.* 2017;26:1770-84.
22. Liu X, Lu D, Saw WY, Shaw PJ, Wangkumhang P, Ngamphiw C, Fucharoen S, Lert-Itthiporn W, Chin-Inmanu K, Chau TN, Anders K, Kasturiratne A, de Silva HJ, Katsuya T, Kimura R, Nabika T, Ohkubo T, Tabara Y, Takeuchi F, Yamamoto K, Yokota M, Mamatyusupu D, Yang W, Chung YJ, Jin L, Hoh BP, Wickremasinghe AR, Ong RH, Khor CC, Dunstan SJ, Simmons C, Tongsima S, Suriyaphol P, Kato N, Xu S, Teo YY. Characterising private and shared signatures of positive selection in 37 Asian populations. *Eur J Hum Genet.* 2017;25:499-508.
23. Kato N, Loh M, Takeuchi F, Verweij N, Wang X, Zhang W, Kelly TN, Saleheen D, Lehne B, Leach IM, Drong AW, Abbott J, Wahl S, Tan ST, Scott WR, Campanella G, Chadeau-Hyam M, Afzal U, Ahluwalia TS, Bonder MJ, Chen P, Dehghan A, Edwards TL, Esko T, Go MJ, Harris SE, Hartiala J, Kasela S, Kasturiratne A, Khor CC, Kleber ME, Li H, Mok ZY, Nakatochi M, Sapari NS, Saxena R, Stewart AF, Stolk L, Tabara Y, Teh AL, Wu Y, Wu JY, Zhang Y, Aits I, Da Silva Couto Alves A, Das S, Dorajoo R, Hopewell JC, Kim YK, Koivula RW, Luan J, Lyytikäinen LP, Nguyen QN, Pereira MA, Postmus I, Raitakari OT, Bryan MS, Scott RA, Sorice R, Tragante V, Traglia M, White J, Yamamoto K, Zhang Y, Adair LS, Ahmed A, Akiyama K, Asif R, Aung T, Barroso I, Bjorntjes A, Braun TR, Cai H, Chang LC, Chen CH, Cheng CY, Chong YS, Collins R, Courtney R, Davies G, Delgado G, Do LD, Doevendans PA, Gansevoort RT, Gao YT, Grammer TB, Grarup N, Grewal J, Gu D, Wander GS, Hartikainen AL, Hazen SL, He J, Heng CK, Hixson JE, Hofman A, Hsu C, Huang W, Husemoen LL, Hwang JY, Ichihara S, Igase M, Isono M, Justesen JM, Katsuya T, Kibriya MG, Kim YJ, Kishimoto M, Koh WP, Kohara K, Kumari M, Kwek K, Lee NR, Lee J, Liao J, Lieb W, Liewald DC, Matsubara T, Matsushita Y, Meitinger T, Mihailov E, Milani L, Mills R, Mononen N, Müller-Nurasyid M, Nabika T, Nakashima E, Ng HK, Nikus K, Nutile T, Ohkubo T, Ohnaka K, Parish S, Paternoster L, Peng H, Peters A, Pham ST, Pinidiyapathirage MJ, Rahman M, Rakugi H, Rolandsson O, Rozario MA, Ruggiero D, Sala CF, Sarju R, Shimokawa K, Snieder

- H, Sparsø T, Spiering W, Starr JM, Stott DJ, Stram DO, Sugiyama T, Szymczak S, Tang WH, Tong L, Trompet S, Turjanmaa V, Ueshima H, Uitterlinden AG, Umemura S, Vaarasmäki M, van Dam RM, van Gilst WH, van Veldhuisen DJ, Viikari JS, Waldenberger M, Wang Y, Wang A, Wilson R, Wong TY, Xiang YB, Yamaguchi S, Ye X, Young RD, Young TL, Yuan JM, Zhou X, Asselbergs FW, Ciullo M, Clarke R, Deloukas P, Franke A, Franks PW, Franks S, Friedlander Y, Gross MD, Guo Z, Hansen T, Jarvelin MR, Jørgensen T, Jukema JW, Kähönen M, Kajio H, Kivimäki M, Lee JY, Lehtimäki T, Linneberg A, Miki T, Pedersen O, Samani NJ, Sørensen TI, Takayanagi R, Toniolo D; BIOS-consortium; CARDIoGRAMplusC4D; LifeLines Cohort Study; InterAct Consortium, Ahsan H, Allayee H, Chen YT, Danesh J, Deary IJ, Franco OH, Franke L, Heijman BT, Holbrook JD, Isaacs A, Kim BJ, Lin X, Liu J, März W, Metspalu A, Mohlke KL, Sanghera DK, Shu XO, van Meurs JB, Vithana E, Wickremasinghe AR, Wijmenga C, Wolffenbuttel BH, Yokota M, Zheng W, Zhu D, Vineis P, Kyrtopoulos SA, Kleinjans JC, McCarthy MI, Soong R, Gieger C, Scott J, Teo YY, He J, Elliott P, Tai ES, van der Harst P, Kooner JS, Chambers JC. Trans-ancestry genome-wide association study identifies 12 genetic loci influencing blood pressure and implicates a role for DNA methylation. *Nat Genet.* 2015;47:1282-93.
24. Saw WY, Liu X, Khor CC, Takeuchi F, Katsuya T, Kimura R, Nabika T, Ohkubo T, Tabara Y, Yamamoto K, Yokota M; Japanese Genome Variation Consortium, Teo YY, Kato N. Mapping the genetic diversity of HLA haplotypes in the Japanese populations. *Sci Rep.* 2015;5:17855.
25. Rodriguez-Araujo G, Nakagami H, Takami Y, Katsuya T, Akasaka H, Saitoh S, Shimamoto K, Morishita R, Rakugi H, Kaneda Y. Low alpha-synuclein levels in the blood are associated with insulin resistance. *Sci Rep.* 2015;5:12081.
26. Koriyama H, Nakagami H, Nakagami F, Osako MK, Kyutoku M, Shimamura M, Kurinami H, Katsuya T, Rakugi H, Morishita R. Long-term reduction of high blood pressure by angiotensin II DNA vaccine in spontaneously hypertensive rats. *Hypertension.* 2015;66:167-74.
27. Takeuchi F, Isono M, Yamamoto K, Yokota M, Akiyama K, Katsuya T, Kim HS, Park JE, Jang Y, Lee JY; AGEN Consortium, Lee JY, Kato N. Heterogeneous effects of association between blood pressure Loci and coronary artery disease in East Asian individuals. *Circ J.* 2015;79:830-8.
28. Hwang JY, Sim X, Wu Y, Liang J, Tabara Y, Hu C, Hara K, Tam CH, Cai Q, Zhao Q, Jee S, Takeuchi F, Go MJ, Ong RT, Ohkubo T, Kim YJ, Zhang R, Yamauchi T, So WY, Long J, Gu D, Lee NR, Kim S, Katsuya T, Oh JH, Liu J, Umemura S, Kim YJ, Jiang F, Maeda S, Chan JC, Lu W, Hixson JE, Adair LS, Jung KJ, Nabika T, Bae JB, Lee MH, Seielstad M, Young TL, Teo YY, Kita Y, Takashima N, Osawa H, Lee SH, Shin MH, Shin DH, Choi BY, Shi J, Gao YT, Xiang YB, Zheng W, Kato N, Yoon M, He J, Shu XO, Ma RC, Kadowaki T, Jia W, Miki T, Qi L, Tai ES, Mohlke KL, Han BG, Cho YS, Kim BJ. Genome-wide association meta-analysis identifies novel variants associated with fasting plasma glucose in East Asians. *Diabetes.* 2015;64:291-8.

29. Congrains A, Kamide K, Hirose N, Arai Y, Oguro R, Nakama C, Imaizumi Y, Kawai T, Kusunoki H, Yamamoto H, Onishi-Takeya M, Takeya Y, Yamamoto K, Sugimoto K, Akasaka H, Saitoh S, Miura T, Awata N, Kato N, Katsuya T, Ikebe K, Gondo Y, Rakugi H. Disease-associated polymorphisms in 9p21 are not associated with extreme longevity. *Geriatr Gerontol Int*. 2014;15:797-803.
30. Akiyama K, Takeuchi F, Isono M, Chakrawarthy S, Nguyen QN, Wen W, Yamamoto K, Katsuya T, Kasturiratne A, Pham ST, Zheng W, Matsushita Y, Kishimoto M, Do LD, Shu XO, Wickremasinghe AR, Kajio H, Kato N. Systematic fine-mapping of association with BMI and type 2 diabetes at the FTO locus by integrating results from multiple ethnic groups. *PLoS One*. 2014 Jun 30;9(6):e101329.
31. Satoh M, Kato N, Hosaka M, Elnagar N, Tsuchihashi T, Yagi N, Ebara F, Uchiba K, Mori H, Kato M, Yamamoto H, Yoshida K, Miyakawa M, Hatori Y, Yamamoto Y, Katsuya T, Watanabe Y, Yatagai S, Sato H, Imai Y. Validity of salt intake assessment system based on a 24-h dietary recall method using a touch panel computer. *Clin Exp Hypertens*. 2014;36:471-7.
32. Kamide K, Asayama K, Katsuya T, Ohkubo T, Hirose T, Inoue R, Metoki H, Kikuya M, Obara T, Hanada H, Thijs L, Kuznetsova T, Noguchi Y, Sugimoto K, Ohishi M, Morimoto S, Nakahashi T, Takiuchi S, Ishimitsu T, Tsuchihashi T, Soma M, Higaki J, Matsuura H, Shinagawa T, Sasaguri T, Miki T, Takeda K, Shimamoto K, Ueno M, Hosomi N, Kato J, Komai N, Kojima S, Sase K, Miyata T, Tomoike H, Kawano Y, Ogihara T, Rakugi H, Staessen JA, Imai Y; GEANE study group; HOMED-BP study group. Genome-wide response to antihypertensive medication using home blood pressure measurements: a pilot study nested within the HOMED-BP study. *Pharmacogenomics*. 2013;14:1709-21.
33. Kelly TN, Takeuchi F, Tabara Y, Edwards TL, Kim YJ, Chen P, Li H, Wu Y, Yang CF, Zhang Y, Gu D, Katsuya T, Ohkubo T, Gao YT, Go MJ, Teo YY, Lu L, Lee NR, Chang LC, Peng H, Zhao Q, Nakashima E, Kita Y, Shu XO, Kim NH, Tai ES, Wang Y, Adair LS, Chen CH, Zhang S, Li C, Nabika T, Umemura S, Cai Q, Cho YS, Wong TY, Zhu J, Wu JY, Gao X, Hixson JE, Cai H, Lee J, Cheng CY, Rao DC, Xiang YB, Cho MC, Han BG, Wang A, Tsai FJ, Mohlke K, Lin X, Ikram MK, Lee JY, Zheng W, Tetsuro M, Kato N, He J. Genome-wide association study meta-analysis reveals transethnic replication of mean arterial and pulse pressure loci. *Hypertension*. 2013;62:853-9.
34. Katsuya T, Morishita R. Gene polymorphism of angiotensin II type 1 and type 2 receptors. *Curr Pharm Des*. 2013;19:2996-3001.
35. Khan TA, Shah T, Prieto D, Zhang W, Price J, Fowkes GR, Cooper J, Talmud PJ, Humphries SE, Sundstrom J, Hubacek JA, Ebrahim S, Lawlor DA, Ben-Shlomo Y, Abdollahi MR, Slooter AJ, Szolnoki Z, Sandhu M, Wareham N, Frikke-Schmidt R, Tybjaerg-Hansen A, Fillenbaum G, Heijmans BT, Katsuya T, Gromadzka G, Singleton A, Ferrucci L, Hardy J, Worrall B, Rich SS, Matarin M, Whittaker J, Gaunt TR, Whincup P, Morris R, Deanfield J, Donald A, Davey Smith G, Kivimaki M, Kumari M, Smeeth L, Khaw KT, Nalls M, Meschia J, Sun K, Hui R, Day I, Hingorani AD, Casas JP. Apolipoprotein e genotype,

cardiovascular biomarkers and risk of stroke: Systematic review and meta-analysis of 14,015 stroke cases and pooled analysis of primary biomarker data from up to 60,883 individuals. *Int J Epidemiol*. 2013;42:475-92.

36. Kyutoku M, Nakagami H, Koriyama H, Nakagami F, Shimamura M, Kurinami H, Tomioka H, Miyake T, Katsuya T, Morishita R. Inhibition of Neointima Formation through DNA Vaccination for Apolipoprotein(a): A New Therapeutic Strategy for Lipoprotein(a). *Sci Rep*. 2013;3:1600
37. Nakagami F, Koriyama H, Nakagami H, Osako MK, Shimamura M, Kyutoku M, Miyake T, Katsuya T, Rakugi H, Morishita R. Decrease in Blood Pressure and Regression of Cardiovascular Complications by Angiotensin II Vaccine in Mice. *PLoS One*. 2013;8:e60493.
38. Ohishi M, Kawai T, Hayashi N, Kitano S, Katsuya T, Nagano M, Hirotsu A, Yamamoto K, Kamide K, Rakugi H. Effect of tablets with a combination of telmisartan and amlodipine on patients with hypertension: the Cotalo study. *Hypertens Res*. 2013;36:620-626.
39. Takeuchi F, Yamamoto K, Isono M, Katsuya T, Akiyama K, Ohnaka K, Rakugi H, Yamori Y, Ogihara T, Takayanagi R, Kato N. Genetic impact on uric acid concentration and hyperuricemia in the Japanese Population. *J Atheroscler Thromb*. 2013;20:351-67.
40. Kawai T, Kamide K, Ito N, Onishi M, Oguro R, Takeya Y, Tatara Y, Maekawa Y, Katsuya T, Ohishi M, Rakugi H. -374 T/A polymorphism in RAGE gene is associated with onset of diabetes mellitus, atherosclerosis, and renal dysfunction in patients with hypertension. *Clin Exp Hypertens*. 2013;35:236-241.
41. Okamura T, Yanobu-Takanashi R, Takeuchi F, Isono M, Akiyama K, Shimizu Y, Goto M, Liang YQ, Yamamoto K, Katsuya T, Fujioka A, Ohnaka K, Takayanagi R, Ogihara T, Yamori Y, Kato N. Deletion of CDKAL1 affects high-fat diet-induced fat accumulation and glucose-stimulated insulin secretion in mice, indicating relevance to diabetes. *PLoS One*. 2012;7:e49055.
42. Takeuchi F, Isono M, Katsuya T, Yokota M, Yamamoto K, Nabika T, Shimokawa K, Nakashima E, Sugiyama T, Rakugi H, Yamaguchi S, Ogihara T, Yamori Y, Kato N. Association of genetic variants influencing lipid levels with coronary artery disease in Japanese individuals. *PLoS One*. 2012;7:e46385.
43. Okada Y, Sim X, Go MJ, Wu JY, Gu D, Takeuchi F, Takahashi A, Maeda S, Tsunoda T, Chen P, Lim SC, Wong TY, Liu J, Young TL, Aung T, Seielstad M, Teo YY, Kim YJ, Lee JY, Han BG, Kang D, Chen CH, Tsai FJ, Chang LC, Fann SJ, Mei H, Rao DC, Hixson JE, Chen S, Katsuya T, Isono M, Ogihara T, Chambers JC, Zhang W, Kooner JS; The KidneyGen Consortium; The CKDGen Consortium, Albrecht E; The GUGC consortium, Yamamoto K, Kubo M, Nakamura Y, Kamatani N, Kato N, He J, Chen YT, Cho YS, Tai ES, Tanaka T. Meta-analysis identifies multiple loci associated with kidney function-related traits in east Asian populations. *Nat Genet*. 2012;44:904-9.

44. Takeuchi F, Yamamoto K, Katsuya T, Sugiyama T, Nabika T, Ohnaka K, Yamaguchi S, Takayanagi R, Ogihara T, Kato N. Reevaluation of the association of seven candidate genes with blood pressure and hypertension: a replication study and meta-analysis with a larger sample size. *Hypertens Res.* 2012;35:825-31.
45. Congrains A, Kamide K, Katsuya T, Yasuda O, Oguro R, Yamamoto K, Ohishi M, Rakugi H. CVD-associated non-coding RNA, ANRIL, modulates expression of atherogenic pathways in VSMC. *Biochem Biophys Res Commun.* 2012;419:612-6.
46. Oguro R, Kamide K, Katsuya T, Akasaka H, Sugimoto K, Congrains A, Arai Y, Hirose N, Saitoh S, Ohishi M, Miura T, Rakugi H. *Exp Gerontol.* A single nucleotide polymorphism of the adenosine deaminase, RNA-specific gene is associated with the serum triglyceride level, abdominal circumference, and serum adiponectin concentration. *Exp Gerontol.* 2012;47:183-7.
47. Congrains A, Kamide K, Oguro R, Yasuda O, Miyata K, Yamamoto E, Kawai T, Kusunoki H, Yamamoto H, Takeya Y, Yamamoto K, Onishi M, Sugimoto K, Katsuya T, Awata N, Ikebe K, Gondo Y, Oike Y, Ohishi M, Rakugi H. Genetic variants at the 9p21 locus contribute to atherosclerosis through modulation of ANRIL and CDKN2A/B. *Atherosclerosis.* 2012;220:449-55.
48. Takeuchi F, Yokota M, Yamamoto K, Nakashima E, Katsuya T, Asano H, Isono M, Nabika T, Sugiyama T, Fujioka A, Awata N, Ohnaka K, Nakatochi M, Kitajima H, Rakugi H, Nakamura J, Ohkubo T, Imai Y, Shimamoto K, Yamori Y, Yamaguchi S, Kobayashi S, Takayanagi R, Ogihara T, Kato N. Genome-wide association study of coronary artery disease in the Japanese. *Eur J Hum Genet.* 2011;20:333-40.
49. Popov S, Silveira A, Wågsäter D, Takemori H, Oguro R, Matsumoto S, Sugimoto K, Kamide K, Hirose T, Satoh M, Metoki H, Kikuya M, Ohkubo T, Katsuya T, Rakugi H, Imai Y, Sanchez F, Leosdottir M, Syvänen AC, Hamsten A, Melander O, Bertorello AM. Salt-inducible kinase 1 influences Na⁺,K⁺-ATPase activity in vascular smooth muscle cells and associates with variations in blood pressure. *J Hypertens.* 2011;29:2395-403.
50. Nakagami H, Shimamura M, Miyake T, Shimosato T, Minobe N, Moritani T, Kiomy Osako M, Nakagami F, Koriyama H, Kyutoku M, Shimizu H, Katsuya T, Morishita R. Nifedipine prevents hepatic fibrosis in a non-alcoholic steatohepatitis. *Mol Med Report.* 2011;5:37-40.
51. Hirose T, Totsune K, Nakashige Y, Metoki H, Kikuya M, Ohkubo T, Hara A, Satoh M, Inoue R, Asayama K, Kondo T, Kamide K, Katsuya T, Ogihara T, Izumi SI, Rakugi H, Takahashi K, Imai Y. Influence of adrenomedullin 2/intermedin gene polymorphism on blood pressure, renal function and silent cerebrovascular lesions in Japanese: the Ohasama study. *Hypertens Res.* 2011;34:1327-32.
52. Nakayama H, Nagai H, Matsumoto K, Oguro R, Sugimoto K, Kamide K, Ohishi M, Katsuya T, Okamoto H, Maeda M, Komamura K, Azuma J, Rakugi H, Fujio Y. Association between osteopontin promoter

variants and diastolic dysfunction in hypertensive heart in the Japanese population. *Hypertens Res.* 2011;34:1141-6.

53. Kato N, Takeuchi F, Tabara Y, Kelly TN, Go MJ, Sim X, Tay WT, Chen CH, Zhang Y, Yamamoto K, Katsuya T, Yokota M, Kim YJ, Ong RT, Nabika T, Gu D, Chang LC, Kokubo Y, Huang W, Ohnaka K, Yamori Y, Nakashima E, Jaquish CE, Lee JY, Seielstad M, Isono M, Hixson JE, Chen YT, Miki T, Zhou X, Sugiyama T, Jeon JP, Liu JJ, Takayanagi R, Kim SS, Aung T, Sung YJ, Zhang X, Wong TY, Han BG, Kobayashi S, Ogihara T, Zhu D, Iwai N, Wu JY, Teo YY, Tai ES, Cho YS, He J. Meta-analysis of genome-wide association studies identifies common variants associated with blood pressure variation in east Asians. *Nat Genet.* 2011;43:531-8.
54. Takeuchi F, Yamamoto K, Katsuya T, Nabika T, Sugiyama T, Fujioka A, Isono M, Ohnaka K, Fujisawa T, Nakashima E, Ikegami H, Nakamura J, Yamori Y, Yamaguchi S, Kobayashi S, Ogihara T, Takayanagi R, Kato N. Association of genetic variants for susceptibility to obesity with type 2 diabetes in Japanese individuals. 2011;54:1350-9.
55. Hirose T, Hashimoto M, Totsune K, Metoki H, Hara A, Satoh M, Kikuya M, Ohkubo T, Asayama K, Kondo T, Kamide K, Katsuya T, Ogihara T, Izumi S, Rakugi H, Takahashi K, Imai Y. Association of (pro)renin receptor gene polymorphisms with lacunar infarction and left ventricular hypertrophy in Japanese women: the Ohasama study. *Hypertens Res.* 2011;34:530-535.
56. Katsuya T, Harrap SB, Ogihara T Genetics of hypertension and cardiovascular disease. *Int J Hypertens.* 2011: 951254.
57. Takeuchi F, Isono M, Nabika T, Katsuya T, Sugiyama T, Yamaguchi S, Kobayashi S, Ogihara T, Yamori Y, Fujioka A, Kato N. Confirmation of ALDH2 as a Major locus of drinking behavior and of its variants regulating multiple metabolic phenotypes in a Japanese population. *Circ J.* 2011;75:911-918.
58. Sugimoto K, Katsuya T, Kamide K, Fujisawa T, Shimaoka I, Ohishi M, Morishita R, Ogihara T, Rakugi H. Promoter Polymorphism of RGS2 Gene Is Associated with Change of Blood Pressure in Subjects with Antihypertensive Treatment: The Azelnidipine and Temocapril in Hypertensive Patients with Type 2 Diabetes Study. *Int J Hypertens.* 2010:196307.
59. Tabara Y, Kohara K, Kita Y, Hirawa N, Katsuya T, Ohkubo T, Hiura Y, Tajima A, Morisaki T, Miyata T, Nakayama T, Takashima N, Nakura J, Kawamoto R, Takahashi N, Hata A, Soma M, Imai Y, Kokubo Y, Okamura T, Tomoike H, Iwai N, Ogihara T, Inoue I, Tokunaga K, Johnson T, Caulfield M, Munroe P, Umemura S, Ueshima H, Miki T. Common Variants in the ATP2B1 Gene Are Associated With Susceptibility to Hypertension. The Japanese Millennium Genome Project. *Hypertension.* 2010;56:973-80.

60. Koriyama H, Nakagami H, Katsuya T, Sugimoto K, Yamashita H, Takami Y, Maeda S, Kubo M, Takahashi A, Nakamura Y, Ogihara T, Rakugi H, Kaneda Y, Morishita R. Identification of Evidence Suggestive of an Association with Peripheral Arterial Disease at the OSBPL10 Locus by Genome-Wide Investigation in the Japanese Population. *J Atheroscler Thromb*. 2010;17:1054-62.
61. Katsuya T, Morishita R. The challenge of stopping the knocking down of metabolic dominos upstream. *Hypertens Res*. 2010;33:529-530.
62. Takeuchi F, Isono M, Katsuya T, Yamamoto K, Yokota M, Sugiyama T, Nabika T, Fujioka A, Ohnaka K, Asano H, Yamori Y, Yamaguchi S, Kobayashi S, Takayanagi R, Ogihara T, Kato N. Blood pressure and hypertension are associated with 7 loci in the Japanese population. *Circulation* 2010;121:2302-2309.
63. Oguro R, Kamide K, Kokubo Y, Shimaoka I, Congrains A, Horio T, Hanada H, Ohishi M, Katsuya T, Okamura T, Miyata T, Kawano Y, Rakugi H. Association of carotid atherosclerosis with genetic polymorphisms of the klotho gene in patients with hypertension. *Geriatr Gerontol Int*. 2010;10:311-8.
64. Michishita T, Kobayashi S, Katsuya T, Ogihara T, Kawabuchi K. Evaluation of the antiobesity effects of an amino acid mixture and conjugated linoleic acid on conjugated overweight humans: a randomized, double-blind, placebo-controlled trial. *J Int. Med. Res*. 2010;38:844-859.
65. Koriyama H, Nakagami H, Katsuya T, Akasaka H, Saitoh S, Shimamoto K, Ogihara T, Kaneda Y, Morishita R, Rakugi H. Variation in OSBPL10 is associated with dyslipidemia. *Hypertens Res*. 2010;33:511-514.
66. Shimaoka I, Kamide K, Ohishi M, Katsuya T, Akasaka H, Saitoh S, Sugimoto K, Oguro R, Congrains A, Fujisawa T, Shimamoto K, Ogihara T, Rakugi H. Association of gene polymorphism of the fat-mass and obesity-associated gene with insulin resistance in Japanese. *Hypertens Res*. 2010;33:214-218.
67. Takeuchi F, Katsuya T, Chakrewarthy S, Yamamoto K, Fujioka A, Serizawa M, Fujisawa T, Nakashima E, Ohnaka K, Ikegami H, Sugiyama T, Nabika T, Kasturiratne A, Yamaguchi S, Kono S, Takayanagi R, Yamori Y, Kobayashi S, Ogihara T, de Silva A, Wickremasinghe R, Kato N. Common variants at the GCK, GCKR, G6PC2-ABCB11 and MTNR1B loci are associated with fasting glucose in two Asian populations. *Diabetologia*. 2010;53:299-308.
68. Watanabe Y, Metoki H, Ohkubo T, Katsuya T, Tabara Y, Kikuya M, Hirose T, Sugimoto K, Asayama K, Inoue R, Hara A, Obara T, Nakura J, Kohara K, Totsune K, Ogihara T, Rakugi H, Miki T, Imai Y. Accumulation of common polymorphisms is associated with development of hypertension: a 12-year follow-up from the Ohasama study. *Hypertens Res*. 2010;33:129-134.
69. Takeuchi F, Nabika T, Isono M, Katsuya T, Sugiyama T, Yamaguchi S, Kobayashi S, Yamori Y, Ogihara T, Kato N. Evaluation of genetic loci influencing adult height in the Japanese population. *J Hum Genet*. 2009;54:749-752.

70. Akasaka H, Katsuya T, Saitoh S, Sugimoto K, Ohnishi H, Congrains A, Ohnishi M, Ohishi M, Rakugi H, Ogihara T, Shimamoto K. A Promoter Polymorphism of Lamin A/C Gene is an Independent Genetic Predisposition to Arterial Stiffness in a Japanese General Population (The Tanno and Sobetsu Study). *J Atheroscler Thromb.* 2009;16:404-409.
71. Miyanaga K, Fukuo K, Akasaka H, Katsuya T, Fukada R, Rakugi H, Kazumi T. C allele of angiotensin II type 1 receptor gene A1166C polymorphism affects plasma adiponectin concentrations in healthy young Japanese women. *Hypertens Res.* 2009;32:901-905.
72. Tomiyama-Hanayama M, Rakugi H, Kohara M, Mima T, Adachi Y, Ohishi M, Katsuya T, Hoshida Y, Aozasa K, Ogihara T, Nishimoto N. Effect of Interleukin-6 Receptor Blockage on Renal Injury in Apolipoprotein E-Deficient Mice. *Am J Physiol Renal Physiol.* 2009;297:F679-684.
73. Katsuya T, Morishita R, Rakugi H, Ogihara T. Genetic basis of hypertension for the development of tailored medicine. *Hypertens Res.* 2009;32:643-648.
74. Hanasaki H, Takemura Y, Fukuo K, Ohishi M, Onishi M, Yasuda O, Katsuya T, Awata N, Kato N, Ogihara T, Rakugi H. Fas promoter region gene polymorphism is associated with an increased risk for myocardial infarction. *Hypertens Res.* 2009;32:261-264.
75. Ueno T, Tabara Y, Fukuda N, Tahira K, Matsumoto T, Kosuge K, Haketa A, Matsumoto K, Sato Y, Nakayama T, Katsuya T, Ogihara T, Makita Y, Hata A, Yamada M, Takahashi N, Hirawa N, Umemura S, Miki T, Soma M. Association of SLC6A9 gene variants with human essential hypertension. *J Atheroscler Thromb.* 2009;16:201-206.
76. Takeuchi F, Serizawa M, Yamamoto K, Fujisawa T, Nakashima E, Ohnaka K, Ikegami H, Sugiyama T, Katsuya T, Miyagishi M, Nakashima N, Nawata H, Nakamura J, Kono S, Takayanagi R, Kato N. Confirmation of multiple risk Loci and genetic impacts by a genome-wide association study of type 2 diabetes in the Japanese population. *Diabetes.* 2009;58:1690-1699.
77. Maekawa Y, Ishikawa K, Yasuda O, Oguro R, Hanasaki H, Kida I, Takemura Y, Ohishi M, Katsuya T, Rakugi H. Klotho suppresses TNF-alpha-induced expression of adhesion molecules in the endothelium and attenuates NF-kappaB activation. *Endocrine.* 2009;35:341-346.
78. Tatara Y, Ohishi M, Yamamoto K, Shiota A, Hayashi N, Iwamoto Y, Takeda M, Takagi T, Katsuya T, Ogihara T, Rakugi H. Macrophage inflammatory protein-1beta induced cell adhesion with increased intracellular reactive oxygen species. *J Mol Cell Cardiol.* 2009;47:104-111.
79. Tsuchihashi-Makaya M, Serizawa M, Yanai K, Katsuya T, Takeuchi F, Fujioka A, Yamori Y, Ogihara T, Kato N. Gene-environmental interaction regarding alcohol-metabolizing enzymes in the Japanese general population. *Hypertens Res.* 2009;32:207-213.

80. Hirose T, Hashimoto M, Totsune K, Metoki H, Asayama K, Kikuya M, Sugimoto K, Katsuya T, Ohkubo T, Hashimoto J, Rakugi H, Takahashi K, Imai Y. Association of (pro)renin receptor gene polymorphism with blood pressure in Japanese men: the Ohasama study. *Am J Hypertens*. 2009;22:294-299.
81. Sugimoto K, Akasaka H, Katsuya T, Node K, Fujisawa T, Shimaoka I, Yasuda O, Ohishi M, Ogihara T, Shimamoto K, Rakugi H. A Polymorphism Regulates CYP4A11 transcriptional activity and is associated with hypertension in a Japanese population. *Hypertension*. 2008;52:1142-8.
82. Katsuya T, Ogihara T. Hypertension and risk stratification. *Hypertens Res* 2008;31:1271-3.
83. Masuo K, Straznicki NE, Lambert GW, Katsuya T, Sugimoto K, Rakugi H, Socratous F, Hastings J, Lambert EA, Ogihara T, Esler MD. Leptin-receptor polymorphisms relate to obesity through blunted leptin-mediated sympathetic nerve activation in a Caucasian male population. *Hypertens Res* 2008;31:1093-1100.
84. Takami Y, Nakagami H, Morishita R, Katsuya T, Hayashi H, Mori M, Koriyama H, Baba Y, Yasuda O, Rakugi H, Ogihara T, Kaneda Y. Potential Role of CYLD (Cylindromatosis) as a Deubiquitinating Enzyme in Vascular Cells. *Am J Pathol*. 2008;172:818-29.
85. Sun X, Yasuda O, Takemura Y, Kawamoto H, Higuchi M, Baba Y, Katsuya T, Fukuo K, Ogihara T, Rakugi H. Akt activation prevents Apop-1-induced death of cells. *Biochem Biophys Res Commun*. 2008;377:1097-101.
86. Kohara K, Tabara Y, Nakura J, Imai Y, Ohkubo T, Hata A, Soma M, Nakayama T, Umemura S, Hirawa N, Ueshima H, Kita Y, Ogihara T, Katsuya T, Takahashi N, Tokunaga K, Miki T. Identification of hypertension-susceptibility genes and pathways by a systemic multiple candidate gene approach: the millennium genome project for hypertension. *Hypertens Res*. 2008;31:203-12.
87. Ueno T, Soma M, Tabara Y, Tokunaga K, Tahira K, Fukuda N, Matsumoto K, Nakayama T, Katsuya T, Ogihara T, Makita Y, Hata A, Yamada M, Takahashi N, Hirawa N, Umemura S, Miki T. Association between Fatty Acid binding protein 3 gene variants and essential hypertension in humans. *Am J Hypertens*. 2008;21:691-695.
88. Mashimo Y, Suzuki Y, Hatori K, Tabara Y, Miki T, Tokunaga K, Katsuya T, Ogihara T, Yamada M, Takahashi N, Makita Y, Nakayama T, Soma M, Hirawa N, Umemura S, Ohkubo T, Imai Y, Hata A. Association of TNFRSF4 gene polymorphisms with essential hypertension. *J Hypertens*. 2008;26:902-13.
89. Kato N, Miyata T, Tabara Y, Katsuya T, Yanai K, Hanada H, Kamide K, Nakura J, Kohara K, Takeuchi F, Mano H, Yasunami M, Kimura A, Kita Y, Ueshima H, 90. Nakayama T, Soma M, Hata A, Fujioka A, Kawano Y, Nakao K, Sekine A, Yoshida T, Nakamura Y, Saruta T, Ogihara T, Sugano S, Miki T, Tomoike

- H. High-density association study and nomination of susceptibility genes for hypertension in the Japanese National Project. *Hum Mol Genet.* 2008;17:617-27
90. Nakagami H, Kikuchi Y, Katsuya T, Morishita R, Akasaka H, Saitoh S, Rakugi H, Kaneda Y, Shimamoto K, Ogihara T. Gene polymorphism of myospryn (cardiomyopathy-associated 5) is associated with left ventricular wall thickness in patients with hypertension. *Hypertens Res.* 2007;30:1239-46.
91. Ohishi M, Takagi T, Ito N, Tatara Y, Hayashi N, Shiota A, Iwamoto Y, Katsuya T, Rakugi H, Ogihara T. Renal protective effect in hypertensive patients -high doses of angiotensin II receptor blocker (HARB) study-. *Hypertens Res* 2007;30:1187-92.
92. Ohishi M, Takagi T, Ito N, Terai M, Tatara Y, Hayashi N, Shiota A, Katsuya T, Rakugi H, Ogihara T. Renal-protective effect of T-and L-type calcium channel blockers in hypertensive patients: an Amlodipine-to-Benidipine Changeover (ABC) study. *Hypertens Res.* 2007;30:797-806.
93. Nakano N, Hori H, Abe M, Shibata H, Arimura T, Sasaoka T, Sawabe M, Chida K, Arai T, Nakahara KI, Kubo T, Sugimoto K, Katsuya T, Ogihara T, Doi Y, Izumi T, Kimura A. Interaction of BMP10 with Tcap may modulate the course of hypertensive cardiac hypertrophy. *Am J Physiol Heart Circ Physiol.* 2007;293:H3396-403.
94. Ch'ng ES, Hoshida Y, Iizuka N, Morii E, Ikeda JI, Yamamoto A, Tomita Y, Hanasaki H, Katsuya T, Maeda K, Ohishi M, Rakugi H, Ogihara T, Aozasa K. Composite malignant pheochromocytoma with malignant peripheral nerve sheath tumour: a case with 28 years of tumour-bearing history. *Histopathology.* 2007;51:420-2.
95. Masuo K, Katsuya T, Sugimoto K, Kawaguchi H, Rakugi H, Ogihara T, Tuck ML. High plasma norepinephrine levels associated with beta2-adrenoceptor polymorphisms predict future renal damage in nonobese normotensive individuals. *Hypertens Res.* 2007;30:503-11.
96. Takami Y, Nakagami H, Morishita R, Katsuya T, Cui TX, Ichikawa T, Saito Y, Hayashi H, Kikuchi Y, Nishikawa T, Baba Y, Yasuda O, Rakugi H, Ogihara T, Kaneda Y. Ubiquitin carboxyl-terminal hydrolase L1, a novel deubiquitinating enzyme in the vasculature, attenuates NF-kappaB activation. *Arterioscler Thromb Vasc Biol.* 2007;27:2184-90.
97. Katsuya T, Rakugi H, Ogihara T. Inflammation and salt sensitivity in the early state of hypertension. *Hypertens Res* 2007;30:105-7.
98. Akasaka H, Katsuya T, Saitoh S, Sugimoto K, Fu Y, Takagi T, Ohnishi H, Rakugi H, Ura N, Shimamoto K, Ogihara T. Effects of angiotensin II type 1 receptor gene polymorphisms on insulin resistance in Japanese general population: the Tannno-Sobetsu Study. *Hypertens Res* 2006;29:961-7.

99. Kawaguchi H, Masuo K, Katsuya T, Sugimoto K, Rakugi H, Ogihara T, Tuck ML. β 2-and β 3-Adrenoceptor polymorphisms relate to subsequent weight gain and blood pressure elevation in obese normotensive individuals. *Hypertens Res* 2006;29:951-9.
100. Yuan M, Ohishi M, Ito N, Sugimoto K, Takagi T, Terai M, Katsuya T, Rakugi H, Wu Z, Ogihara T. Genetic influences of β -adrenoceptor polymorphisms on arterial functional changes and cardiac remodeling in hypertensive patients. *Hypertens Res* 2006;29:875-81
101. Masuo K, Katsuya T, Kawaguchi H, Fu Y, Rakugi H, Ogihara T, Tuck ML. Beta2-adrenoceptor polymorphisms relate to obesity through blunted leptin-mediated sympathetic activation. *Am J Hypertens*. 2006;19:1084-91.
102. Nakayama T, Kuroi N, Sano M, Tabara Y, Katsuya T, Ogihara T, Makita Y, Hata A, Yamada M, Takahashi N, Hirawa N, Umemura S, Miki T, Soma M. Mutation of the follicle-stimulating hormone receptor gene 5' -untranslated region associated with female hypertension. *Hypertension*. 2006;48:512-8.
103. Yamamoto K, Ohishi M, Katsuya T, Ito N, Ikushima M, Kaibe M, Tatara Y, Shiota A, Sugano S, Takeda S, Rakugi H, Ogihara T. Deletion of angiotensin-converting enzyme 2 accelerates pressure overload-induced cardiac dysfunction by increasing local angiotensin II. *Hypertension*. 2006;47:718-26.
104. Xu R, Ishikawa K, Katsuya T, Fu Y, Akasaka H, Ikushima M, Chihara Y, Ura N, Shimamoto K, Rakugi H, Ogihara T. Association between senescence-related uncoupling protein 2 gene polymorphisms and abdominal obesity in Japanese subjects: the Tanno and Sobetsu Study. *Geriatr Gerontol Int*. 2006;6:15-9.
105. Masuo K, Katsuya T, Kawaguchi H, Fu Y, Rakugi H, Ogihara T, Tuck ML. Rebound weight gain as associated with high plasma norepinephrine levels that are mediated through polymorphisms in the beta2-adrenoceptor. *Am J Hypertens*. 2005;18:1508-1516
106. Masuo K, Katsuya T, Fu Y, et al. β 2- and β 3-adrenergic receptor polymorphisms are related to the onset of weight gain and blood pressure elevation over 5 Years. *Circulation*. 2005;111:3429-4.
107. Masuo K, Katsuya T, Fu Y, Rakugi H, Ogihara T, Tuck ML. Beta2-adrenoceptor polymorphisms relate to insulin resistance and sympathetic overactivity as early markers of metabolic disease in nonobese, normotensive individuals. *Am J Hypertens*. 2005;18:1009-1014.
108. Iwashima Y, Katsuya T, Ishikawa K, et al. Association of hypoadiponectinemia with smoking habit in men. *Hypertension*. 2005;45:1094-1100.
109. Masuo K, Katsuya T, Fu Y, et al. Lys418Asn polymorphism of the α 2-adrenoceptor gene relates to serum uric acid levels but not to insulin sensitivity. *Hypertension* 2005;46:144-150.

110. Masuo K, Katsuya T, Ogihara T, Tuck ML. Acute hyperinsulinemia reduces plasma leptin levels in insulin-sensitive Japanese men. *Am J Hypertens.* 2005;18:235-43..
111. Fu Y, Katsuya T, Matsuo A, Yamamoto K, Akasaka H, Takami Y, Iwashima Y, Sugimoto K, Ishikawa K, Ohishi M, Rakugi H, Ogihara T. Relationship of bradykinin B2 receptor gene polymorphism with essential hypertension and left ventricular hypertrophy. *Hypertens Res.* 2004;27:933-8
112. Matsuo A, Katsuya T, Ishikawa K, Sugimoto K, Iwashima Y, Yamamoto K, Ohishi M, Rakugi H, Ogihara T. G2736A polymorphism of thiazide-sensitive Na-Cl cotransporter gene predisposes to hypertension in young women. *J Hypertens.* 2004;22:2123-7.
113. Sugimoto K, Katsuya T, Ohkubo T, Hozawa A, Yamamoto K, Matsuo A, Rakugi H, Tsuji I, Imai Y, Ogihara T. Association between angiotensin II type 1 receptor gene polymorphism and essential hypertension: the Ohasama Study. *Hypertens Res.* 2004;27:551-6.
114. Iwashima Y, Katsuya T, Ishikawa K, Ouchi N, Ohishi M, Sugimoto K, Fu Y, Motone M, Yamamoto K, Matsuo A, Ohashi K, Kihara S, Funahashi T, Rakugi H, Matsuzawa Y, Ogihara T. Hypoadiponectinemia is an independent risk factor for hypertension. *Hypertension.* 2004;43:1318-23.
115. Motone M, Katsuya T, Ishikawa K, Iwashima Y, Sugimoto K, Yamamoto K, Fu Y, Matsuo A, Ohishi M, Rakugi H, Ogihara T. Association between hepatocyte growth factor gene polymorphism and essential hypertension. *Hypertens Res.* 2004;27:247-51.
116. Fukuchi K, Katsuya T, Sugimoto K, Kuremura M, Kim HD, Li L, Ogihara T. LMNA mutation in a 45 year old Japanese subject with Hutchinson-Gilford progeria syndrome. *J Med Genet.* 2004;41:e67.
117. Ji Q, Ikegami H, Fujisawa T, Kawabata Y, Ono M, Nishino M, Ohishi M, Katsuya T, Rakugi H, Ogihara T. A common polymorphism of uncoupling protein 2 gene is associated with hypertension. *J Hypertens.* 2004;22:97-102.
118. Ogihara T, Katsuya T, Ishikawa K, Matsuo A, Rakugi H, Shoji M, Yasujima M. Hypertension in a patient with Gitelman's syndrome. *J Hum Hypertens.* 2004;8:677-9.
119. Ohashi K, Ouchi N, Kihara S, Funahashi T, Nakamura T, Sumitsuji S, Kawamoto T, Matsumoto S, Nagaretani H, Kumada M, Okamoto Y, Nishizawa H, Kishida K, Maeda N, Hiraoka H, Iwashima Y, Ishikawa K, Ohishi M, Katsuya T, Rakugi H, Ogihara T, Matsuzawa Y. Adiponectin I164T mutation is associated with the metabolic syndrome and coronary artery disease. *J Am Coll Cardiol.* 2004;43:1195-200.
120. Takiuchi S, Mannami T, Miyata T, Kamide K, Tanaka C, Kokubo Y, Koyama Y, Inamoto N, Katsuya T, Iwai N, Kawano Y, Ogihara T, Tomoike H. Identification of 21 single nucleotide polymorphisms in

human hepatocyte growth factor gene and association with blood pressure and carotid atherosclerosis in the Japanese population. *Atherosclerosis*. 2004;173:301-7.

- 121.Katsuya T, Sugimoto K, Hozawa A, Ohkubo T, Yamamoto K, Matsuo A, Ishikawa K, Matsubara M, Rakugi H, Tsuji I, Imai Y, Ogihara T. Genetic risk factors for cerebral infarction using data from a large scale genetic epidemiological study: the Ohasama Study. *Geriatr Gerontol Int*. 2003;3:150-3.
- 122.Katsuya T, Ogihara T. Future medicine in the post-genome age -Targeting tailored medicine for hypertension-. *JMAJ*. 2003;46:327-32.
- 123.Katsuya T, Ishikawa K, Sugimoto K, Rakugi H, Ogihara T. Salt sensitivity of Japanese from the viewpoint of gene polymorphism. *Hypertens Res*.2003;26:521-5.
- 124.Inamoto N, Katsuya T, Kokubo Y, Mannami T, Asai T, Baba S, Ogata J, Tomoike H, Ogihara T. Association of methylenetetrahydrofolate reductase gene polymorphism with carotid atherosclerosis depending on smoking status in a Japanese general population. *Stroke*. 2003;34:1628-33.
- 125.Matsubara M, Metoki H, Katsuya T, Kikuya M, Suzuki M, Michimata M, Araki T, Hozawa A, Tsuji I, Ogihara T, Imai Y. T+31C polymorphism (M235T) of the angiotensinogen gene and home blood pressure in the Japanese general population: the Ohasama Study. *Hypertens Res*. 2003;26:47-52.
- 126.Kikuya M, Sugimoto K, Katsuya T, Suzuki M, Sato T, Funahashi J, Katoh R, Kazama I, Michimata M, Araki T, Hozawa A, Tsuji I, Ogihara T, Yanagisawa T, Imai Y, Matsubara M. A/C1166 gene polymorphism of the angiotensin II type 1 receptor (AT1) and ambulatory blood pressure: the Ohasama Study. *Hypertens Res*. 2003;26:141-5.
- 127.Sugimoto K, Hozawa A, Katsuya T, Matsubara M, Ohkubo T, Tsuji I, Motone M, Higaki J, Hisamachi S, Imai Y, Ogihara T. alpha-Adducin Gly460Trp polymorphism is associated with low renin hypertension in younger subjects in the Ohasama study. *J Hypertens*. 2002;20:1779-84.
- 128.Shintani M, Ikegami H, Fujisawa T, Kawaguchi Y, Ohishi M, Katsuya T, Higaki J, Shimamoto K, Ogihara T. Leptin gene polymorphism is associated with hypertension independent of obesity. *J Clin Endocrinol Metab*. 2002;87:2909-12.
- 129.Okuda T, Sumiya T, Mizutani K, Tago N, Miyata T, Tanabe T, Kato H, Katsuya T, Higaki J, Ogihara T, Tsujita Y, Iwai N, Muzutani K. Analyses of differential gene expression in genetic hypertensive rats by microarray. *Hypertens Res*. 2002;25:249-55.
- 130.Morimoto S, Okaishi K, Onishi M, Katsuya T, Yang J, Okuro M, Sakurai S, Onishi T, Ogihara T. Deletion allele of the angiotensin-converting enzyme gene as a risk factor for pneumonia in elderly patients. *Am J Med*. 2002;112:89-94.

131. Mizutani K, Sugimoto K, Okuda T, Katsuya T, Miyata T, Tanabe T, Higaki J, Ogihara T, Yamori Y, Tsujita Y, Tago N, Iwai N. Kynureninase is a novel candidate gene for hypertension in spontaneously hypertensive rats. *Hypertens Res.* 2002;25:135-40.
132. Matsubara M, Suzuki M, Fujiwara T, Kikuya M, Metoki H, Michimata M, Araki T, Kazama I, Satoh T, Hashimoto J, Hozawa A, Ohkubo T, Tsuji I, Katsuya T, Higaki J, Ogihara T, Satoh H, Imai Y. Angiotensin-converting enzyme I/D polymorphism and hypertension: the Ohasama study. *J Hypertens.* 2002;20:1121-6.
133. Komai N, Ohishi M, Moriguchi A, Yanagitani Y, Jinno T, Matsumoto K, Katsuya T, Rakugi H, Higaki J, Ogihara T. Low-dose doxazosin improved aortic stiffness and endothelial dysfunction as measured by noninvasive evaluation. *Hypertens Res.* 2002;25:5-10.
134. Katsuya T, Baba S, Ishikawa K, Mannami T, Fu Y, Inamoto N, Asai T, Fukuda M, Higaki J, Ogata J, Ogihara T. Epsilon 4 allele of apolipoprotein E gene associates with lower blood pressure in young Japanese subjects: the Suita Study. *J Hypertens.* 2002;20:2017-21.
135. Iwai N, Katsuya T, Mannami T, Higaki J, Ogihara T, Kokame K, Ogata J, Baba S. Association between SAH, an acyl-CoA synthetase gene, and hypertriglyceridemia, obesity, and hypertension. *Circulation.* 2002;105:41-7.
136. Fukuda M, Ohkubo T, Katsuya T, Hozawa A, Asai T, Matsubara M, Kitaoka H, Tsuji I, Araki T, Satoh H, Higaki J, Hisamichi S, Imai Y, Ogihara T. Association of a mast cell chymase gene variant with HDL cholesterol, but not with blood pressure in the Ohasama study. *Hypertens Res.* 2002;25:179-84.
137. Fujiwara T, Katsuya T, Matsubara M, Mikami T, Ishikawa K, Kikuya M, Ohkubo T, Hozawa A, Michimata M, Suzuki M, Metoki H, Asayama K, Araki T, Tsuji I, Higaki J, Satoh H, Hisamichi S, Ogihara T, Imai Y. T+31C polymorphism of angiotensinogen gene and nocturnal blood pressure decline: the Ohasama study. *Am J Hypertens.* 2002;15:628-32.
138. Tsujita Y, Iwai N, Katsuya T, Higaki J, Ogihara T, Tamaki S, Kinoshita M, Mannami T, Ogata J, Baba S. Lack of association between genetic polymorphism of CYP11B2 and hypertension in Japanese: the Suita Study. *Hypertens Res.* 2001;24:105-9.
139. Tsujita Y, Baba S, Yamauchi R, Mannami T, Kinoshita M, Yamamoto R, Katsuya T, Higaki J, Ogihara T, Ogata J, Iwai N. Association analyses between genetic polymorphisms of endothelial nitric oxide synthase gene and hypertension in Japanese: The Suita Study. *J Hypertens.* 2001;19:1941-8.
140. Takagi S, Baba S, Iwai N, Fukuda M, Katsuya T, Higaki J, Mannami T, Ogata J, Goto Y, Ogihara T. The aldehyde dehydrogenase 2 gene is a risk factor for hypertension in Japanese but does not alter the sensitivity to pressor effects of alcohol: the Suita study. **Hypertens Res.** 2001;24:365-70.

141. Matsubara M, Kikuya M, Ohkubo T, Metoki H, Omori F, Fujiwara T, Suzuki M, Michimata M, Hozawa A, **Katsuya T**, Higaki J, Tsuji I, Araki T, Ogihara T, Satoh H, Hisamichi S, Nagai K, Kitaoka H, Imai Y. Aldosterone synthase gene (CYP11B2) C-334T polymorphism, ambulatory blood pressure and nocturnal decline in blood pressure in the general Japanese population: the Ohasama Study. **J Hypertens**. 2001;19:2179-84.
142. Mannami T, **Katsuya T**, Baba S, Inamoto N, Ishikawa K, Higaki J, Ogihara T, Ogata J. Low potentiality of angiotensin-converting enzyme gene insertion/deletion polymorphism as a useful predictive marker for carotid atherogenesis in a large general population of a Japanese city: the Suita study. **Stroke**. 2001;32:1250-6.
143. **Katsuya T**, Ogihara T. Current genetics of essential hypertension. **Curr Hypertens Rep**. 2001;3:1-2.
144. **Katsuya T**, Iwashima Y, Sugimoto K, Motone M, Asai T, Fukuda M, Fu Y, Hatanaka Y, Ohishi M, Rakugi H, Higaki J, Ogihara T. Effects of antihypertensive drugs and gene variants in the renin-angiotensin system. **Hypertens Res**. 2001;24:463-7.
145. Iwai N, Baba S, Mannami T, **Katsuya T**, Higaki J, Ogihara T, Ogata J. Association of sodium channel gamma-subunit promoter variant with blood pressure. **Hypertension**. 2001;38:86-9.
146. Ishikawa K, Baba S, **Katsuya T**, Iwai N, Asai T, Fukuda M, Takiuchi S, Fu Y, Mannami T, Ogata J, Higaki J, Ogihara T. T+31C polymorphism of angiotensinogen gene and essential hypertension. **Hypertension**. 2001;37:281-5.
147. Higaki J, **Katsuya T**, Morishita R, Ogihara T. Symposium on the etiology of hypertension—summarizing studies in 20th century. 1. Hypertension and genes. **Intern Med**. 2001;40:144-7.
148. Fu Y, **Katsuya T**, Higaki J, Asai T, Fukuda M, Takiuchi S, Hatanaka Y, Rakugi H, Ogihara T. A common mutation of low-density lipoprotein receptor gene is associated with essential hypertension among Japanese. **J Hum Hypertens**. 2001;15:125-30.
149. Fu Y, **Katsuya T**, Asai T, Fukuda M, Inamoto N, Iwashima Y, Sugimoto K, Rakugi H, Higaki J, Ogihara T. Lack of correlation between Mbo I restriction fragment length polymorphism of renin gene and essential hypertension in Japanese. **Hypertens Res**. 2001;24:295-8.
150. Asai T, Ohkubo T, **Katsuya T**, Higaki J, Fu Y, Fukuda M, Hozawa A, Matsubara M, Kitaoka H, Tsuji I, Araki T, Satoh H, Hisamichi S, Imai Y, Ogihara T. Endothelin-1 gene variant associates with blood pressure in obese Japanese subjects: the Ohasama Study. **Hypertension**. 2001;38:1321-4.
151. Takami S, Imai Y, **Katsuya T**, Ohkubo T, Tsuji I, Nagai K, Satoh H, Hisamichi S, Higaki J, Ogihara T. Gene polymorphism of the renin-angiotensin system associates with risk for lacunar infarction. The Ohasama study. **Am J Hypertens**. 2000;13:121-7.

152. Sato N, **Katsuya T**, Nakagawa T, Ishikawa K, Fu Y, Asai T, Fukuda M, Suzuki F, Nakamura Y, Higaki J, Ogihara T. Nine polymorphisms of angiotensinogen gene in the susceptibility to essential hypertension. **Life Sci.** 2000;68:259-72.
153. Ohishi M, Rakugi H, Miki T, **Katsuya T**, Okamura A, Kamide K, Nakata Y, Takami S, Ikegami H, Yanagitani Y, Tabuchi Y, Kumahara Y, Higaki J, Ogihara T. Deletion polymorphism of angiotensin-converting enzyme gene is associated with postprandial hyperglycaemia in individuals undergoing general check-up. **Clin Exp Pharmacol Physiol.** 2000;27:483-7.
154. Ogihara T, **Katsuya T**, Higaki J. Genetic analysis of essential hypertension in Japanese populations. **Ann N Y Acad Sci.** 2000;902:8-16.
155. Matsubara M, Ohkubo T, Michimata M, Hozawa A, Ishikawa K, **Katsuya T**, Nagai K, Tsuji I, Higaki J, Araki T, Satoh H, Hisamichi S, Ito S, Ogihara T, Imai Y. Japanese individuals do not harbor the T594M mutation but do have the P592S mutation in the C-terminus of the beta-subunit of the epithelial sodium channel: the Ohasama study. **J Hypertens.** 2000;18:861-6.
156. **Katsuya T**, Sato N, Asai T, Fukuda M, Takiuchi S, Fu Y, Hanayama R, Moriguchi A, Rakugi H, Higaki J, Ogihara T. Association of polymorphism in the promoter region of the apolipoprotein E gene with diastolic blood pressure in normotensive Japanese. **Hypertens Res.** 2000;23:271-5.
157. Ishikawa K, Imai Y, **Katsuya T**, Ohkubo T, Tsuji I, Nagai K, Takami S, Nakata Y, Satoh H, Hisamichi S, Higaki J, Ogihara T. Human G-protein beta3 subunit variant is associated with serum potassium and total cholesterol levels but not with blood pressure. **Am J Hypertens.** 2000;13:140-5.
158. Higaki J, Baba S, **Katsuya T**, Sato N, Ishikawa K, Mannami T, Ogata J, Ogihara T. Deletion allele of angiotensin-converting enzyme gene increases risk of essential hypertension in Japanese men : the Suita Study. **Circulation.** 2000;101:2060-5.
159. Okamura A, Ohishi M, Rakugi H, **Katsuya T**, Yanagitani Y, Takiuchi S, Taniyama Y, Moriguchi K, Ito H, Higashino Y, Fujii K, Higaki J, Ogihara T. Pharmacogenetic analysis of the effect of angiotensin-converting enzyme inhibitor on restenosis after percutaneous transluminal coronary angioplasty. **Angiology.** 1999;50:811-22.
160. Iwai N, **Katsuya T**, Ishikawa K, Mannami T, Ogata J, Higaki J, Ogihara T, Tanabe T, Baba S. Human prostacyclin synthase gene and hypertension : the Suita Study. **Circulation.** 1999;100:2231-6.
161. Takemoto Y, Sakatani M, Takami S, Tachibana T, Higaki J, Ogihara T, Miki T, **Katsuya T**, Tsuchiyama T, Yoshida A, Yu H, Tanio Y, Ueda E. Association between angiotensin II receptor gene polymorphism and serum angiotensin converting enzyme (SACE) activity in patients with sarcoidosis. **Thorax.** 1998;53:459-62.

162. Takami S, Katsuya T, Rakugi H, Sato N, Nakata Y, Kamitani A, Miki T, Higaki J, Ogihara T. Angiotensin II type 1 receptor gene polymorphism is associated with increase of left ventricular mass but not with hypertension. **Am J Hypertens**. 1998;11:316-21.
163. Nakata Y, Katsuya T, Takami S, Sato N, Fu Y, Ishikawa K, Takiuchi S, Rakugi H, Miki T, Higaki J, Ogihara T. Methylenetetrahydrofolate reductase gene polymorphism: relation to blood pressure and cerebrovascular disease. **Am J Hypertens**. 1998;11:1019-23.
164. Nakamura S, Moriguchi A, Morishita R, Aoki M, Yo Y, Hayashi S, Nakano N, Katsuya T, Nakata S, Takami S, Matsumoto K, Nakamura T, Higaki J, Ogihara T. A novel vascular modulator, hepatocyte growth factor (HGF), as a potential index of the severity of hypertension. **Biochem Biophys Res Commun**. 1998;242:238-43.
165. Ishikawa K, Katsuya T, Sato N, Nakata Y, Takami S, Takiuchi S, Fu Y, Higaki J, Ogihara T. No association between alpha-adducin 460 polymorphism and essential hypertension in a Japanese population. **Am J Hypertens**. 1998;11:502-6.
166. Sato N, Katsuya T, Rakugi H, Takami S, Nakata Y, Miki T, Higaki J, Ogihara T. Association of variants in critical core promoter element of angiotensinogen gene with increased risk of essential hypertension in Japanese. **Hypertension**. 1997;30:321-5.
167. Nakata Y, Katsuya T, Rakugi H, Takami S, Sato N, Kamide K, Ohishi M, Miki T, Higaki J, Ogihara T. Polymorphism of angiotensin converting enzyme, angiotensinogen, and apolipoprotein E genes in a Japanese population with cerebrovascular disease. **Am J Hypertens**. 1997;10:1391-5.
168. Katsuya T, Horiuchi M, Minami S, Koike G, Santoro NF, Hsueh AJ, Dzau VJ. Genomic organization and polymorphism of human angiotensin II type 2 receptor: no evidence for its gene mutation in two families of human premature ovarian failure syndrome. **Mol Cell Endocrinol**. 1997;127:221-8.
169. Kamide K, Rakugi H, Nakano N, Ohishi M, Nakata Y, Takami S, Katsuya T, Higaki J, Ogihara T. Insulin resistance is related to silent cerebral infarction in patients with essential hypertension. **Am J Hypertens**. 1997;10:1245-9.
170. Haywood GA, Gullestad L, Katsuya T, Hutchinson HG, Pratt RE, Horiuchi M, Fowler MB. AT1 and AT2 angiotensin receptor gene expression in human heart failure. **Circulation**. 1997;95:1201-6.
171. Wu DA, Bu X, Warden CH, Shen DD, Jeng CY, Sheu WH, Fuh MM, Katsuya T, Dzau VJ, Reaven GM, Lusis AJ, Rotter JI, Chen YD. Quantitative trait locus mapping of human blood pressure to a genetic region at or near the lipoprotein lipase gene locus on chromosome 8p22. **J Clin Invest**. 1996;97:2111-8.

172. Takami S, Higaki J, Miki T, Katsuya T, Nakata Y, Rakugi H, Serikawa T, Ogihara T. Analysis and comparison of new candidate loci for hypertension between genetic hypertensive rat strains. **Hypertens Res.** 1996;19:51-6.
173. Nakata Y, Katsuya T, Rakugi H, Takami S, Ohishi M, Kamino K, Higaki J, Tabuchi Y, Kumahara Y, Miki T, Ogihara T. Polymorphism of the apolipoprotein E and angiotensin-converting enzyme genes in Japanese subjects with silent myocardial ischemia. **Hypertension.** 1996;27:1205-9.
174. Katsuya T, Takami S, Higaki J, Serikawa T, Mikami H, Miki T, Ogihara T. Gap junction protein locus on chromosome 18 cosegregates with body weight in the spontaneously hypertensive rat. **Hypertens Res.** 1995;18:63-7.
175. Katsuya T, Horiuchi M, Chen YD, Koike G, Pratt RE, Dzau VJ, Reaven GM. Relations between deletion polymorphism of the angiotensin-converting enzyme gene and insulin resistance, glucose intolerance, hyperinsulinemia, and dyslipidemia. **Arterioscler Thromb Vasc Biol.** 1995;15:779-82.
176. Katsuya T, Koike G, Yee TW, Sharpe N, Jackson R, Norton R, Horiuchi M, Pratt RE, Dzau VJ, MacMahon S. Association of angiotensinogen gene T235 variant with increased risk of coronary heart disease. **Lancet.** 1995;345:1600-3.
177. Kamino K, Nagano K, Katsuya T, Nishiwaki Y, Takeda M, Tanabe H, Nishimura T, Ii K, Fujimoto K, Tsujimura R, et al. Linkage and haplotype analysis of familial early-onset Alzheimer disease in Japanese population. **Jpn J Hum Genet.** 1995;40:229-41.
178. Morishita R, Rakugi H, Higaki J, Tomita N, Nakamura F, Yu H, Katsuya T, Mikami H, Ogihara T. Differential regulation of brain angiotensin II in genetically hypertensive and normotensive rats after nephrectomy. **Blood Press.** 1994;3:265-9.
179. Mullan M, Tsuji S, Miki T, Katsuya T, Naruse S, Kaneko K, Shimizu T, Kojima T, Nakano I, Ogihara T, et al. Clinical comparison of Alzheimer's disease in pedigrees with the codon 717 Val→Ile mutation in the amyloid precursor protein gene. **Neurobiol Aging.** 1993;14:407-19.
180. Katsuya T, Higaki J, Zhao Y, Miki T, Mikami H, Serikawa T, Ogihara T. A neuropeptide Y locus on chromosome 4 cosegregates with blood pressure in the spontaneously hypertensive rat. **Biochem Biophys Res Commun.** 1993;192:261-7.
181. Higashimori K, Zhao Y, Higaki J, Kamitani A, Katsuya T, Nakura J, Miki T, Mikami H, Ogihara T. Association analysis of a polymorphism of the angiotensin converting enzyme gene with essential hypertension in the Japanese population. **Biochem Biophys Res Commun.** 1993;191:399-404.

182. Katsuya T, Higaki J, Miki T, Kohara K, Yagisawa H, Tanase H, Mikami H, Serikawa T, Nojima H, Ogihara T. Hypotensive effect associated with a phospholipase C-delta 1 gene mutation in the spontaneously hypertensive rat. **Biochem Biophys Res Commun.** 1992;187:1359-66.
183. Yoshioka K, Miki T, Katsuya T, Ogihara T, Sakaki Y. The 717Val—Ile substitution in amyloid precursor protein is associated with familial Alzheimer's disease regardless of ethnic groups. **Biochem Biophys Res Commun.** 1991;178:1141-6.
184. Katsuya T, Higaki J, Miki T, Nakura J, Ikegami H, Morishita R, Nagano M, Higashimori K, Nakamura F, Mikami H, et al. Substrain comparison of genetically hypertensive rats using DNA fingerprinting, and genetic analysis of blood pressure in the inbred rats. **Tohoku J Exp Med.** 1991;165:253-60